

The image shows the front cover of a book. The cover is primarily a light-colored, textured material, possibly cloth or paper, with a dark red, textured material (likely leather or faux leather) covering the spine and the corners. The cover is decorated with a series of concentric rectangular borders. The outermost border is a wide band with a repeating pattern of small circles. Inside this is a narrower band with a repeating pattern of small diamonds. The central area of the cover is divided into two horizontal sections by a decorative horizontal line with a scalloped or wavy pattern. The word "RECORD" is printed in a bold, serif, all-caps font in the center of the upper section. The spine of the book is visible on the left, showing the same light-colored material and some wear at the top and bottom edges.

RECORD

SHOUT
SHARE IT



Collection and Field Note Book

卷之四

(March 10, 1952 - December 16, 1952)

(20448 - 51448)

(No. 1 to 46, G. I. Gravelly Coll.)

(Continued from Book No. 35)

100

.01

Bokehok lalat, dua lalat, seekor lalat.

Enclaves: latest, Ujane latest, Wotja

.....

Užtas laiškas 1944-71443.....

88-56 SS443, te[al nie[al ewX, [lo3A nie[al ewX

General Remarks 34-50

(Faint, illegible text)

77-25 10442-25442, 10442-25442, 10442-25442
10442-25442, 10442-25442, 10442-25442

[illegible]

.....

.....

CO-50 26742-10742 5514 Newborn
22-53 5514/5 4/12/11 011111

.....

.....

.....

.....

25-26

.....

10-77 ***** 2000-00-00
30 60 ***** 10/000000

.....

.....

[illegible][illegible]

Collection and Field Note Book

No. 33

(March 10, 1952 - December 16, 1952)

(34417 - 34496)

- (No. 1 to 46, G. L. Gressitt Coll.) -

(Continued from Book No. 32)

	<u>Page No.</u>
MARSHALL ISLANDS	
Ujae Atoll, Bik Islet, Alle (Erlia) Islet, Bikenkar Islet, Enelamoj Islet, Bokerok Islet, Rua Islet, Bock Islet, Enelamoj Islet, Ujae Islet, Wojia Islet, Bock Islet.....	1-23
Ujae Islet 34417-34421	24-31
Kwajalein Atoll, Kwajalein Islet, 34422	32-33
General Remarks	34-50
Wotho Atoll, Wotho Islet, 34423-34424	56-57
Enejelto Islet 34425-34426	58
Enearik Islet 34427-34429	58
Ombelim Islet 34430	60-61
Kabben Islet 34431-34432	62-63
Ombelin Islet 34433	62-63
Biken Islet 34434	62-63
Kabben Islet 34435-34436	64-67
Eneobnak Islet..... 34437-34443	68-71
Enebarbar Islet	72-73
Mejurwor Islet	72-76
Eneobnak Islet	77-81
Bokanaetok	82-85
Wotho Islet	85-86
Bokanaetok	86
Iroi jemau Islet	86-87

	<u>Page No.</u>
MARSHALL ISLANDS	
Wotho Atoll, Enejelto Islet	88-89
Kabben Islet (ctd. from p. 65).....	90-98
Enearik Islet	99-101
Wotho birds	102-108
Itinerary of voyage - from Capt. M. W. Shoap's log book of FS 367.....	110-111
Bikar Atoll, Jolliklik Islet, Almeni Islet, Jaboero Islet, Bikar Islet.....	112-114
Birds	115
Pokak (Taongi Atoll, birds	116-120
Kwajalein Atoll	121
MARIANAS ISLANDS	
Guam	122-125
Uracas	125
VOLCANO ISLAND	
Iwo Jima 34444-34445	126-127
JAPAN	
Chichibu Range	128-129
WAKE ISLAND ATOLL	130-135
..... continued	142-148
Wake Islet 34446-34483	136-143
Birds	149-152
VIRGINIA, Fairfax 34484-34489	156-157
" " 34491-34496	160-161
MARYLAND, Anne Arundel Co. 34490	158-159
- KAYARGEL ATOLL (G.L. Gressitt Coll.).....	
Ngariungs Islet..... 1-27	166-167
Ngajaugel Islet 28-43b	168-169
- PALAU, Auluptagel Island near Koror 44-46 .	170-171

End.

begin 34417
end 34496

F. R. Fosberg

Collection book #33
+ field data



Standard Miniature Blank Book

No. 665 9½ x 6 120 Pages Units
No. 667 9½ x 6 200 Pages Units
No. 668 9½ x 6 300 Pages Units

Made in the Following Rulings

Journals, Day or Cash Books, Double \$ and Cts.

S. E. Ledgers, \$ and Cts.

Double Entry Ledgers

Records with Margin Line

When ordering give Number and Ruling desired

Made in U. S. A.

WHEN ORDERING THIS BOOK SPECIFY
STOCK NUMBER AND RULING DESIRED.

A BOORUM & PEASE PRODUCT

1952 Marshall Is. Ujae Atoll 1

3

h. 162

(Continued from Book No. 32)

March 10 - Bar somewhat north of Bik Islet - gravel bar, submerged at ~~low~~ high tide. Not held, as far as can be seen from lagoon, by any beach rock or other stable material. Seems to represent a balance between the strength of the lagoon waves and that of the sea waves weakened by their passage across a wide reef flat. At almost high tide there were small breakers on both sides of it.

At rather low tide the bar was well exposed and hundreds of noddies were holding a conference on it.

nt
yllus.
sis
ll,
her
h
r
tin-
ch,
t
her-
he
lge
ies
ing
sage
are
ll
ees,
end,
ser-
ola
rowth
ndergrowth.
ravel
teep
on
his
le

March 10 - Alby (Erlie) Islet
 Three patches of coconuts, one near south lagoon point, two near center of islet.

Center of islet occupied by *Pisonia* grove ^{12-15 m. high} - small grove of four large trees and half-dozen or so smaller ones, open beneath with only a few scattered *Morinda* and *Allophylus* ~~seed~~ saplings, almost no ground cover - a few scattered small plants of *Thuarea*, *Polypodium*, and *Ipomoea tuba* (seedlings). Ground mostly bare and center covered by a layer of 2-6" of peat. (see p. 186-7 of Book 32 for description of this, sample #110).

This grove is on a very slight rise in the ground.

Surrounding it is mixed open forest of *Pisonia*, *Pandanus*, *Terminalia* and two of the patches of coconuts.

Undergrowth and ground cover similar to that in *Pisonia* grove but a little more dense, same composition but with addition of *Boerhavia tetrandra* and ~~the~~

seedlings of the dominant trees, much more *Allophylus*. One *Terminalia samoensis* tree is about 10 m. tall, with 3 main branches from near base each 3 dm. thick.

Lagoon side rather sandy, with one continuous ^{low} ridge along beach, a second back of it interrupted by a depression in middle. The n. end of this second ridge corresponds with a piece of lagoonward dipping beach-rock on the passage beach. The ridges are covered by rather tall but sparse ^{mixed} scrub forest, with a ^{small} patch of young coconut trees, 5-6 m. tall on ~~so~~ south end, one or two on north, *Messerschmidia* and *Scaevola* dominant, much *Xylocarpus* in undergrowth. ^{Weedy in openings + tangled in undergrowth.} South passage beach, except for ~~the~~ gravel hook on lagoon end, a steep boulder beach up from reef-rock, on top of this a beach and weak ridge of pebble to cobble

size ~~rocks~~ stones. Along this passage a scrub forest of *Messerschmidia* and *Pisonia*, interrupted by patches of *Scaevola* scrub ~~and~~ with scattered *Messerschmidia* trees taller than the scrub.

Seaward apex of islet and outer 50 m. a flat of broken coral at height of passage ridges. In from this is rubble getting finer toward the lagoon. Along the north passage is a cobble ridge, weak at seaward end, stronger lagoonward. The outer half of islet, except in ~~very~~ center, is a sparse scrub or scrub-forest 3-5 m. high of *Messerschmidia*, *Scaevola*, *Pisonia*, *Quettarda*, with *Wedelia* tangled in it and in openings, ground cover of *Boerhaavia* and *Polypodium*.

Lagoonward half of north passage beach lined by almost pure stand of *Cordia*, with some *Pisonia*, between coconut patch and beach

and extending in both directions. Absolutely no undergrowth or ground cover. Canopy very sparse at this season.

Hermit crabs very abundant on this islet, in day-time hiding in and under rotten coconut logs, trash, etc. Large series collected (Jan 13, 14, 15) to see how many species represented. Color patterns vary considerably. Of large red species, color varies from scarlet to brick-red to brownish-red.

March 9 - ^{Bikini} Biginnigan Islet
 A flat triangle of sand and fine rubble coarser toward the windward periphery where there is a boulder beach and, on the south-east side, at least, a thickly scattered boulders along the edge. Center rather sandy. Leeward beach a broad sandy beach prolonged into two horns at the corners, these curving to leeward.

Vegetation a sparse mixture of *Messerschmidia*, *Scaevola*, and *Guettarda*, a few *Pisonia*, *Intsia* scrub-forest to leeward, becoming lower to windward. Much open ground between bushes to windward this covered by *Lepturus*, *Cassytha*, *Triumfetta*, *Boerhavia tetrandra*, *Polypodium*. Through the central portion are scattered coconut trees, these very yellow and unhealthy-looking, some of them dead.

Leeward corner, on photo

almost bare, now quite well vegetated with *Scaevola* and a few *Messerschmidia*, some of the latter of good size (3 m. ±).

Judging by the photos and present observations the island is adding sand on its inner side and the sand horns at the corners change shape very rapidly.

(Chief Ehti's father told Max Neil that during the strong trade-wind months these ^{horns on all islets} bend strongly inward, while during June to September they straighten out parallel with beach. This should be checked with photos of M.I. project.)

1952 Marshall Is.

Enlamoj

Mar. 9 - Enlamoj Islet

On south half of leeward reef flat are scattered large boulders.

Greater part of island planted to coconuts, fairly closely spaced, mostly healthy looking, except for one spot toward seaward side about $\frac{1}{4}$ from north end, where the trees are yellow, some have died, even the seedlings are very yellow (shows as a light spot on photo M1-38-25).

Soil generally sand to fine ~~to~~ loam, locally fine rubble, gray to gray brown.

A few areas have been cleared out recently and have a ground-cover of *Trumfetta*, *Fimbristylis*, *Polypodium*, etc. Large areas with mat of *Wedelia* 1-2 m. thick. Others with thickets of *Guettarda*, *Morinda*, *Bougainvillea*, etc.

One patch of *Intsia* near n.w. corner, and scattered in plantation generally.

General second story of *Pandanus*, scattered to quite frequent, locally in very sparse spots in plantation *Pandanus*.

Ujae atoll

could be called dominant.

Locally in north ~~on~~ half, near center, (dark trees in M1-38-25, 26) are patches of seedy var. of breadfruit. No undergrowth or ground cover here except scattering of breadfruit seedlings up to 3-4 dm. tall.

Here and there patches of mixed forest, *Pisonia*, and *Achrosia*, *Intsia*, ^{*Pandanus*}, etc., and, especially toward south end, patches of *Achrosia* forest, some of these,

toward ~~outer~~ windward beach, very yellow ^(light on photo), some trees dead. ^(near beach)

Fuzzy-looking trees south of light spot on photo M1-38-25 are *Pisonia*.

Leeward beach is sand and fine gravel. At top is an interrupted narrow belt of *Scaevola* scrub.

Beach around north west coast very broad, sandy, no significant seaward ridge. Around on north ~~west~~ side a sand ridge is well developed but tapers off southward. This coast lined with broad belt of tall scrub or scrub forest of *Scaevola*,

Messerschmidia, guettarda, etc., tangled with Wedelia, Cassytha, etc. Soil here very sandy. Scrub immediately at top of n.e. beach relatively low, mostly Scaevola. This scrub extends southward along entire east beach, becoming taller, scrub forest on boulder ridge.

North-East beach is of boulders except about 200 m. at each end, which is sand. The boulder ridge is low, perhaps ~~20~~ 30 m. wide, covered by scrub forest described above, which extends inward for some distance with addition of ^{Pandanus} Pisonia, Ochrosia, etc. Tongues of it extend well into coconut plantation. Here and there, in mixed forest near beach are small groups of very wind-beaten and bedraggled coconut trees.

Along ^{east} ~~west~~ side of ^{south and} are small dunes. Beach here is sand. Scrub belt narrow here. Much mixed forest in plantation.

boulders thickly scattered 30 m. more inward, but with much sand around them.

on seaward side.

South end a sand flat with sparse Scaevola scrub, much bare sand.

y. 145

Birds seen on Ujae Atoll.

According to Chief Enti, all sea-birds are eaten, young or old, but heron is not eaten, as it eats rats.

MacNeil reports seeing a noddy (which he not determined) pick up a veritid snail from reef, fly high in air, drop it, following it down, so its shell broke on rocks and it could be eaten. This accounts for numerous freshly broken veritid shells on rocks (sample # III from Rock I, Z. coll. MacNeil).

"aali"* Frigate birds (*Fregatta minor*) A large colony of at least 75 birds live on Pokairuk Islet, south of Boek Channel, where they may be seen roosting in some numbers in *Pisonia* trees almost any time. At sunset they are soaring in great numbers over the channel. A few of them have no white at all, either beneath or on heads.

Brown booby - "kalo"
About 30 seen in evening (Mar. 1) returning to Pokairuk Islet after day's fishing. One seen on Ujae I. (arrow) one on Biginnigan

Red-footed booby -
One seen dead on outer beach of Rua I. on Feb. 23.

"aali"* Common noddy - "piguak"
At least occasionally seen on all islets, some seen gathering nesting material in forest on Boek I.; hundreds seen on Pokairuk I.; quite a few on Rua I. ^{Feb. 24} three nests in *Terminalia* tree on seaward side. The nests may be distinguished from those of the white-capped noddy by being as much as twice as large, of looser construction, made largely of twigs, not

much stained with guano. On Wotjis Islet, over the lagoon beach, six noddies were seen vigorously pursuing a fairy tern that had a small fish in its beak. The chase lasted several minutes, but in spite of the wavering, fluttering type of flight of the fairy tern it got away from its apparently sure flying assailants.

On Biginbigan Islet, at low tide, ^{March 9} an enormous flock of hundreds or even thousands of noddies blackened a long sand-spit uncovered at low tide. They were holding the usual "conference" so characteristic of common noddies. In the afternoon, at high tide many were flying over the islet. One nest with an egg in it was found in a bush about 2 m. above the ground. The egg was white with brown markings. The old bird protested and flew around close by, trying to drive me off.

Many were seen flying about Enyharu Islet, and they were abundant on ally I. Here one was nesting in a big crotch of a Pisonia tree about 5 m. above the ground. ~~the~~ The nest in this broad depressed crotch consisted of very little material, a few twigs and leaves, not at all typical for this species.

On Ujae islet only six noddies seen. This islet is where the people are concentrated. (Mar. 13)

White-capped Noddy -
Feb. 16-21 "jagarik" *
Nesting abundantly and gregariously in Pisonia trees on Bok and Pokairuk islets, and to be seen in hundreds flying around these islets. Young fully feathered but not yet flying. Natives collect these young in considerable numbers.

to eat. They built a fire at base of a huge *Pisonia* loaded with nests. It burned all day and finally one half of the tree fell. at least ~~at least~~ five or six dozen young were collected. called "nemajelij kon".

Common on ally Islet, a dozen nesting in close group in *Pisonia* and *Terminalia* trees, Mar. 10.

Many nesting gregariously on Rua I. Feb. 23. The nests may be distinguished from those of common noddy ~~because~~ because they are smaller, more compact, built mostly of leaves (*Pisonia* and *Messerschmidia*), badly stained with white guano. Usually found in groups.

Fairy tern "mejan" *
seen occasionally at any hour of the day carrying ~~a~~ small fish in their beaks.

Many in trees and flying overhead on Boek, one young one in pinfeather stage, Feb. 17.

Hundreds flying overhead on Polkairuk.

Hundreds also seen on Rua, Feb. 23, six eggs seen, three on *Messerschmidia*, three on *Pisonia*, one broken one under *Pisonia*, two young almost feathered out.

egg
7 mm.
long.

Only occasionally seen on Ebejer Islet, rare on Ujae.

Hundreds seen over Bile, many on ~~the~~ Enklamij.

Abundant on ally Islet, one nesting on with egg on *Pisonia* branch at about 8 m., drove off noddy when it approached.

One ~~over~~ over lagoon near Ebejer with fish in mouth out flew and eluded six common noddies that were pursuing it.

Black-naped tern

One seen flying over south end of Boak I. Feb. 19.

Seven on Rura, on inner south sand-spit of islet ^{Feb. 19}. Two nests here on bare gravel flat at base of sand spit, just above high tide level. Eggs laid on absolutely bare gravel, 2 in one clutch, 1 in other, very well concealed by resemblance to surroundings. Eggs identical in size, shape, color and markings with those of *Gygis alba* - gray ground color with scattered brownish mottling, 29 mm. long.

Four seen on inner southwest sand spit of ~~Boak~~ Ally Islet, Mar. 10; one clutch of two eggs laid on small collection of pebbles on bare sand, ^{just above high tide,} very well concealed, only the frantic protests of the birds calling my attention to the eggs even though I had been walking around them for some minutes. Looked for them carefully before

locating them. They were exactly as described above. As long as I was in vicinity, even all during time I ate my lunch sitting on rock 50' away, the four birds protested noisily, "yakh, yakh, yakh" "gip, eep, eep" "aa" "yakh, yakh, gip" etc.

Crested tern - "ke'er" (name tied to this species by description by natives only, as no bird seen when native was present).

On Mar. 8, two seen over lagoon, Ebej I.

On Mar. 9, three seen on ^{Pikidun} Bigin Nigan and one over lagoon in north part.

On Mar. 10 eight seen over seaward reef flat, Ally Islet.

On Mar. 13, one seen over lagoon near windward reef about 2 mi. n. of Ujae I.

Pacific Golden Plover
- "kolej"*

Not as common here
as on some atolls.

Two seen on reef flat
at north end of Bokl I.

One seen on Rua I.

On Ebeju one was seen
on lagoon beach in flock
of turnstones, ~~also~~ ^{in flock on}
seaward beach.

Four in flock with
turnstones and curlews
on Wotjia I. on reef flat.

Two seen on north end
of Enyglamij I.

Four in flock with
curlew on Ally I.

An Ujae one seen on
south beach and one on
east end beach.

Wandering Tattler - "kijij"

Most often seen on
exposed reef flats or rubble
flats, ~~also~~ usually on
seaward side of islets.

On Bokl I. four were
seen in a flock on the outer
reef flat, two together on
rubble flat on lagoon side.

One to three seen on each
of Ujae, Rua, Wotjia, Ebeju,

Ally, ^{Ujae} ~~Enyglamij~~ ^{Enyglamij}, and
Enyglamij islets.

Note, usually sounded
when they fly from one
spot to another, is a high
"dee-dee-dee-dee-dee". They
usually fly low over
reef or sea just outside
it. On Enyglamij seen
in flock with turnstones.

Turnstone "kutkut" or
"kutekut".

Five in flock seen on
outer reef, north end of Bokl I.

Flock of 7-8 on Ujae (Arrow),
flock of 5 on outer reef (Torrey).

Four seen on Rua I., one of
these in grassy opening
in interior.

Flock of 10 on ~~seaward~~
lagoon beach, one of 5 on
seaward reef flat, Ebeju I.

Flock of ~~seven~~ 7 on outer
reef flat with plovers and
curlews, Wotjia I.

Flocks of 14 and 7 on
north and northeast
beaches of Enyglamij I.

Flock of 9 on passage
beach, Bijiinigan I.

Bristle-thighed curlew -
"kowak"

Considerable variation
in amount of chestnut
in plumage.

On Boak Islet 1-3 & be
seen almost any time on
different sides of islet,
probably same ones, are
seen in rocky opening in
Scaevola scrub.

4-5 on Ruda Islet, seen
several times.

2 seen on seaward reef of
Ebeja I.

2 on Wotjia (with plovers
and turnstones).

1 seen near south end on
east side of Enyhamij I.

5 seen together on south
passage beach, Biginnigar I.

1 seen with plovers, three
seen ~~in~~ ^{together} together, on
Ally I.

Large droppings frequently
seen on beach filled with
Scaevola stones must
belong to this bird.

Note when flying "kiwee,
kiwee"

Whimbrel

Less common than
Bristle-thighed curlew.

One seen on Ujae by arrow,
one by me (identified by
note rather than color).

One on Ruda I., not
certain if it was this
species.

Two seen on seaward
reef. Ebeja

Note when flying
"dee-dee-dee-dee-dee". This
is often best means
of distinguishing this
species from the Bristle-
thighed curlew.

Reef Heron - "kawij" or "kavij"

Only white and dark
phases seen, all occurrences
listed below, though some may
be repeated flushings on different
days of same bird.

White	Dark
Boak 1	Boak 1
Ujae 2	Ujae 3
Ebeja 2	Ebeja 4
Wotjia 1	Enyhamij 1
Bik 1	
Biginnigar 1	

March 13 - Ujae Islet -

34417 *Artocarpus altilis* (Park.) Fosb.
common mixed with
coconuts in interior
of islet

18 *Premna obtusifolia* R. Br.
one tree seen in village

19 *Allophylus timorensis* (D.C.) Blume
~~common~~ occasional in scrub
along passage (~~common~~ ^{occasional}
also in plantation)

5 20

on coral boulders in
scrub on seaward side

21 *Pseuderanthemum* ^{*canthioides* var.} *atropurpureum*
cultivated in village

#112 - drift seeds from Ujae

#113 - drift seeds from Ailuk

Vial #133 - ants from large
colony in rotten coconut
trunk, Ujae Islet, Mar. 13.
sting painfully. Also a
small millipede from same
coconut trunk

Vial #134 - ants and associated
scale insects - scales on
panicle and calyx of
coconuts, ants running
up and down tree, tending them.
beetle flying.

tree 15 m. tall, 5 dm thick,
buttressed at base;
leaves variable in
outline.

small tree, sterile.

small tree 5 m. tall,
flowers white.

(Bull.) ^{Fosb.} ~~Smiley~~

shrub, 1 m. tall; leaves
dark bronze-purple.
flowers purplish pink,
thickly dotted with red.

Marshallese word
for *Tridacna gigas* -
"kaburi"

Mar. 13 - Ujae Islet.

On seaward side entire beach is of pebbles & cobbles steep, its crest forming a small seaward ridge. This beach and ridge hook around on both ends, to half or two thirds the width of the islet.

Back of the beach is a very broad low ridge gradually and unevenly fading off inward to small rubble and soil about 150 - 200 m. in from beach. The remainder of the islet has an unevenly flat surface of fine rubble or rubble and gray or blackish loam, mixed, this resting on rubble a few inches down.

The island is mostly planted to coconuts, mixed here and there with breadfruit, very locally with *Pisonia*. The second story is of scattered *Pandanus*, locally *Guettarda*, with *Papayas* quite common beneath. Some bananas, but these looking markedly yellow, much more so than when we were here a month ago. *Allophyllus* and *Morinda*.



dunes

DEPARTMENT OF THE INTERIOR
UNITED STATES
GEOLOGICAL SURVEY
WASHINGTON, D. C.

uneven boundary between coarse and fine rubble
rubble ridge

pebble-cobble beach

Plantation

on fine
rubble soil
(mixed with
loam, locally)

cemented
Boulders
on reef flat

over

000 - 100 11 - 0 - 0.4



UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY
WASHINGTON 25, D. C.

when we were here a year
ago. *Allophylus* and *Morinda*,

especially *Morinda*, are
also common. Locally
mats of *Wedelia* up to
a meter deep occur in
the plantation. Degree of
clearing out the brush
varies from place to place.
Young coconuts are very
thick in places. *Vigna*,
~~the~~ *Polypodium*, *Lepturus*,
Thuarea, etc. form ground
cover, any one of which
may be dominant locally,
the *Vigna* especially so
near outer edge of islet.
Inner and passage edges of
island lined with a narrow
belt of scrub or scrub forest.
Along the ~~west~~ ^{west} passage
beach this is composed of
Scaevola and *Guettarda*,
with some *Messerschmidia*
and *Messerschmidia*
and *Morinda*, with a
little *Terminalia*, *Allophylus*
and *Pandanus*, and,
away from beach, *Pisonia*.
Ground cover here of *Lepturus*
or *Vigna* or a mixture
of both, or with *Trumpfetta*
and *Thuarea*. Some
Wedelia tangled in
this belt.

Around the west end and looking around the western part of the seaward beach is a curved, outward dipping bed of beach-~~rock~~ conglomerate, much eroded and pitted even discontinuous in place. 10-20 m. out from present beach, forming a low cuesta. Possibly indicates a former shore-line.

The western end of the seaward beach is coarse gravel with a very broad (150 m. (?) rubble ridge behind it. The outer edge of this, for varying widths, is covered by almost pure *Scaevola* scrub, with a little *Quettarda* and *Messerschmidia*.

Inside this, the rest of the rubble ridge is cleared and planted to coconuts. The ground cover here is of *Vigna* with ~~Lepturus~~ *Lepturus*, *Wedelia*, and *Polypodium*. In places soil is accumulated on top of ridge. The surface is very uneven.

Eastward the outer ridge becomes sharper and steeper and the broad area we become higher.

Further in
Lepturus
becomes
dominant
and *Tacca*,
Polypodium
& *Thunbergia*
common

In the outer ^{edge} ~~half~~ of the plantation *Scaevola* is a common small tree.

On the southeast coast the rubble changes to large boulders and the outer edge of the plantation gives way to *Prisonia* - *Quettarda* forest with a few scattered coconuts.

Many seedling coconuts in undergrowth which is a tangled mass of *Wedelia* and *Ipomoea tuba*.

A little further along this is again replaced by scrub or scrub forest, mostly *Scaevola*. at edge, mixed with *Quettarda* and *Messerschmidia* further back. Locally at edge there is a clump of large *Messerschmidia* trees. Here the pebble-cobble beach is very steep, rather undercut, but with few roots exposed even though cut back almost to bases of trees.

On the reef flat here boulders are very thickly scattered.

On the east end these boulders are piled together and partially cemented into conglomerate on the smooth reef-flat surface. This seems quite recent.

The rubble ridge narrows rapidly north of middle of east end, and ends at same point where cobble beach gives way to fine gravel. The scrub belt narrows and becomes a scrub forest of *Scaevola*, *Messerschmidia* and *Guettarda*. There is much *Pisonia* and breadfruit and some *Ochrosia* in the east end of the plantation, as well as *Morinda*, *Allophylus* and *Pandanus* in second story. Locally breadfruit ~~is~~ is dominant. Papayas are abundant. Ground cover mainly of *Thuarea* and *Vigna*.

The interior generally ~~is~~ has a gently rolling topography.

On the inner west corner

are low sand dunes.

March 15 - Kwajalein Islet
1 white reef heron seen on
seaward reef flat.

Plovers common generally.
3 turnstones seen flying
over island.

Beach rock beds at northern
end of islet on lagoon side
and in passage dipping
away from islet.

34422 green algae
in beach rock just below
high tide level.

The dry spell has
continued. The grass
does not look much
worse than at beginning
of month, but some
Fimbristylis seems
almost dead.

Coconut and *Messerschmidia*
trees transplanted from
marine compound to
swimming pool, with
quite small balls of
earth on roots, are nearly
dead, probably will not
recover, at least unless
there is rain.

(1) Probable component communities of original vegetation:

(a) Relatively stable a "climax" community:

1. *Pisonia* forest -

Tall forest, 20-30 m. high, canopy very dense except in dry seasons when trees may become more or less bare. ^{no} undergrowth ^{except root sprouts,} and usually no ground cover. Tendency to form peat on surface of ground. Usually a pure stand or almost so. Very widespread but relatively small stands.

2. *Ochrosia* forest -

Tall forest, 20-25 m. high or sometimes ^{not quite so} tall, ^{pure stands,} canopy very dense, no undergrowth except thick stands of *Ochrosia* seedlings which seldom attain more than 0.5 m. except under thin places in canopy. Peculiar tessellate appearance on aerial photos. Rather widespread ⁱⁿ small to large stands.

3 *Barringtonia* forest -

Tall forest, 25 m. very large trees, pure stand, canopy complete, no undergrowth or ground cover whatever. Tessellate appearance on aerial photos, difficult to tell from *Ochrosia*, but finer grained. Only one stand known (Lae I. Lae Atoll)

3a *Intsia bijuga* forest

3b *Condis subcordata* forest

~~3c *Intsia bijuga* forest~~

4 *Pemphis* forest -

Rather low forest - 5-7 m., dense, pure stand, usually dense lower branches, often dead, make penetration difficult and painful, usually on exposed rock, varies to a scrub in exposed localities, tolerates pure salt water over roots at high tides. A soft dull green appearance from a distance. Small stands but widespread.

5. Mixed forest -

A variable mixture, basically Messerschmidia, Quettarda, and Pandanus, with usual addition of one or more of Pisonia, Intsia, Ochrosia, ^{Alphitoea}, Cordia, Terminalia, ^{Premna}, and Bougainvillea. May vary in proportions and any of the secondary species may locally assume dominance, occasionally even form pure stands locally.

This is usually tangled with Ipomoea tuba, often with Wedelia, often has undergrowth of seedlings of tree species. In open places may have ground cover

Fimbristylis, of Lepturus, or Thunbergia, or Polypodium, or Trumbetta, or a mixture of the ~~two~~ ^{first} five.

The canopy varies from loose to complete. This varies to scrub-~~scrub~~ forest or scrub, with addi-

tion of Scaevola, Suriana, Pemphis, etc. Scaevola scrub is an extreme of this but so commonly encountered that it should be treated separately.

- (2) Instability is a normal characteristic of atoll habitats - storms, typhoons, tsunamis, etc., altering it by removal or deposit of material. Therefore stable communities may be expected to be relatively unusual.

(b) Pioneer communities:

1. Lepturus grassland - Pure stand of bushy or loose mat-like Lepturus, with some admixture locally of other herbaceous species. Subject to invasion by woody species and by Cassytha. Persistence in places like Pokah (if the opening, there are of this ~~off~~ type) or on small dry windward islets must

be due to very dry conditions.

2. *Fimbristylis grassland* -
Pure stands of *Fimbristylis*
cymosa - occur usually
on dry rocky substrata.
May be mixed with
other herbaceous species,
~~especially *Thespesia*~~
and tends to give
way to woody types.

3. *Scaevola* - *Messerschmidia*
scrub -

Sparsely to dense
mixture of these
two components,
with occasional
addition of *Pemphis*,
Terminalia, *Suriana*
and *Guettarda*. This
is an extreme form
of the mixed forest
community existing
under very pioneer
conditions and possibly
developing into the
latter. It is the
characteristic early
vegetation on sand
bars, sand spits,
denuded areas, etc.

4. *Scaevola* scrub -

an extreme form of
no. 3, commonly of
pure *Scaevola frutescens*,
found as a fringe
along almost all
seaward beaches,
sometimes on lagoon
beaches. On ~~the~~
windward coast,
it may form a
rather broad belt,
sloping to windward,
giving way on
inner side to mixed
scrub or scrub forest.

(3) Successional phenomena.

1. *Messerschmidia* tends not to reproduce itself in well-developed mixed forest.

2. The mixed forest - basically *Messerschmidia*, *Guettarda*-*Pandanus*, is, in more mesophytic habitats, a temporary community, in which the secondary components may, singly or in combination, eventually dominate.

3. The seedlings of *Ochrosia*, *Pandanus*, *Soulaea* and root sprouts of *Pisonia* tolerate their own shade and that of other trees very well. These, then, should under certain circumstances, be able to form relatively pure stable communities, and do. *Intsia*, also, can do this to some extent.

4. *Ochrosia* can seed

itself so effectively and does so well in the shade that it is able to crowd out other seedlings, and, in the general Kwajalein latitude, at least, the presence of a healthy tree of *Ochrosia* means that the forest it is in is doomed to replacement by *Ochrosia*.

5. *Allophyllus* seedlings can get started if there is a thin place in the canopy, and have been observed invading ~~the~~ *Ochrosia* forest in the places where the *Ochrosia* is yellow and unhealthy.

6. *Scaevola* and sometimes *Guettarda*, very successfully invade *Lepturus* grassland.

7. *Wedelia* can also invade *Lepturus* and may form a dense mat a meter or more deep which is invaded only with difficulty.

8. *Scaevola*, *Messerschmidia*,
Quettarda, *Lepturus*,
Triumfetta. *Vigna*
 commonly appear as
 seedlings on ~~very~~
~~salt~~ open sand or
 gravel, on beaches,
 bars, flats, spits, etc.
 where they are exposed
 to pure salt water.

9 *Cassytha* seems to
 be able to invade
 and parasitize
 anywhere except
 in deep shade.

(4) Processes occurring in atoll soils:

1. Deposition of miscellaneous calcareous materials by storm waves.
2. Addition of calcareous fine material or (lime) sand by wind. This commonly deposited around vegetation.
3. Addition of floating pumice.
4. Weathering of pumice.
5. Gradual solution of calcareous material.
5. Removal of material by wind.
6. Removal of material by storm waves.
7. Addition of sea salts ~~in~~ from spray and storm waves.
8. Binding of surface particles by
8. Addition of N, P, Ca, etc. ~~to~~ in guano by fish-eating birds.

9. Binding of surface particles by crust-forming blue-green algae.
10. Binding of loose materials by vegetation.
11. Percolation - seeping downward of rain-water, taking dissolved substances with it.
12. Capillary rise of water, bringing with it any dissolved material in top layer of free water lens, also any picked up on the way up.
13. Nitrogen-fixation by crust-forming blue-green algae (?)
14. Nitrogen-fixation by Rhizobium in root nodules of Vigna, Canavalia, and possibly other legumes.
15. Addition of humus from roots in ground.
16. Addition of humus from leaves, twigs, branches, and trunks on surface of the ground.

- 17 Retention of mineral matter in form not so readily leached out in organic material - plant and animal bodies, humus, etc.
- 18 Enrichment of mineral content of soil by ~~poor~~ weathering products of pumice.
- 19 Oxidation of organic matter exposed to air
- 20 Breaking down of organic matter by bacteria and fungi.
- 21 Breaking down of organic matter by crabs, millipeds, isopods, insects, snails and earthworms.
- 22 Bringing of unaltered materials to surface by burrowing land crabs.
- 23 Evaporation of water from surface of soil.
- 24 Extraction of water by roots.
- 25 Extraction of minerals by roots.

- 26 Formation of *Pisonia* peat.
- 27 Solution of Ca^{P}H_2 by peat acids.
- 28 Solution of Ca^{P}CO_3 by guano acids.
- 29 Cementing of sand under peat and guano.
- 30 Reduction of guano increment by frightening off or destruction of sea-birds.
- 31 Composting of taro pits.
- 32 Muck-formation in bottoms of taro pits.
- 33 Stirring up of surface layers by chickens and hogs.
- 34 Removal of P in copra exported.
- 35 Burning of coconut trash and other dead plant materials by natives.
- 36 Reduction of soil fauna and flora by this burning.
- 37 Loss of humus by burning.

- 38 Consequent loss of fixed N.
 39 Compaction of soil by traffic.
 40 Partial cementation of exposed layers.

41 Binding of loose material by vegetation.

45 Binding of mineral matter in form not so readily washed out by rain in organic material - plant & animal bodies, humus, etc

42 Compaction of soil by traffic.

43 Partial cementation of exposed layers.

44 Oxidation of organic matter exposed to air.

19 Evaporation of water from surface.

20 Extraction of water ~~in~~ by roots.

47 Extraction of minerals by roots.

48 Addition of ^{and supporting} floating particles.

47 Binding of loose material by vegetation.

48 Binding of mineral matter in form not so readily washed out by rain in organic material - plant & animal bodies, humus, etc.

49 Compaction of soil by traffic.

50 Partial cementation of exposed layers.

51 Oxidation of organic matter exposed to air.

52 Evaporation of water from surface.

53 Extraction of water ~~and~~ by roots.

54 Extraction of minerals by roots.

55 Addition of ^{and weathering to} floating particles.

- 118 Breaching down of surface by crabs, gophers, field mice, etc. and plants that have roots in soil in old logs, etc.
- 122 Formation of tundra peats.
- 123 Discoloring of water by peat water.
- 124 Cementing of sand, mud, silt, etc. and gravel.
- 130 Burrowing of animals, tracks, etc. of other kinds of plants, material by nature.
- 131 Accumulation of soil, formation of humus by the building.
- 139 Removal of phosphate in copra.
- 140 Reduction of phosphate content when it is used for fertilizers, etc. or reduced in numbers.
- 142 Composting of tar pits.
- 147 Much formation in bottoms of tar pits.
- 148 Stirring up of surface layers by chickens and dogs.

42 Addition of guano - N & P. by
gull-eating birds.

43 Burrowing in all soil materials
by burrowing land crabs.

44 Burrowing in the soil by
small crustaceans, e.g. amphipods,
etc. - green algae.

45 Nitrogen fixation by crust-forming
blue-green algae. ?

46 Nitrogen fixation by Rhizobia in
roots of Vigna, Canavalia, etc.

47 Regeneration of organic matter
by bacteria and fungi.

Soil processes

- ✓8 Percolation - movement of water through soil by gravity, distribution of water in soil at various subsoils.
- ✓9 Capillary rise of water, bringing water up from deeper layers, and its evaporation in to large open free water, clouds.
- ✓10 Addition of ^{miss.} calcareous material by wind. This commonly ~~deposited~~ deposits around vegetation.
- ✓11 Deposition of ^{miss.} calcareous material by storm waves.
- ✓12 Removal of material by wind.
- ✓13 Removal of material by storm waves.
- ✓15 Addition of salt from spray and waves.
- ✓13 Addition of land mass from rocks in ground.
- ✓14 Addition of ^{miss.} limestone from ^{miss.} limestone, ^{miss.} sand, ^{miss.} shells, and ^{miss.} other ^{miss.} pieces of ground.

50

1952 Marshall Is.

38 Consequent loss of fixed N.

39 Compaction of soil by traffic.

40 Partial cementation of exposed layers.

general remarks

51

52

1952 Marshall Is.

General remarks

53

54

1952

Marshall Is.

General remarks

55

March 13 - Wotho Islet -

Vial 135 - bottom layer -
beetles found eating
holes in overripe fallen
breadfruit (Wotho? Mar. 18).

34423

3

in surface layer of
beach sandstone between
tides

3

24

on surface of base of
old coconut root crown
(sample 114)

sample #114 - pieces of
peat-like remains of old
coconut root crown - for
comparison with samples
#105 & 106, to be sure that there
are not also remains of such
old root crowns.

Along inside of windward
reef, south of Wotho
Islet, and particularly
south of the long sand
islet south of it, were
some of the "oil streaks"
described from Ujae. Most
seemed quite broken up
by wind, but several were
very strong. Some were

#135 (old) top layer - fly, mosquito
and cricket found in forest,
Kabben I. Mar. 20.

green layer.

black

noticed inside of open
reef, some inside the
sand islet. There
were nowhere as
many as commonly
Ujae.

March 19 - ^{English} Enickerudo Islet

34425

green algae

3

discolored beach conglomerate,
about high tide level on
seaward side, south end.

3

26 Auricularia

on dead Ochrosia trunk

Vial 136 - insects and
spiders - orb weavers in
bush and woods; yellow ants
and black ants from two
colonies in dead Messerschmidia
~~stem~~ trunk, each in separate
stub, but within 1 dm. ~~from~~
larvae, etc. go with the black ones.

March 19 - ^{English} Eirek Islet

4

27 Cordia subcordata Lam.

large spreading, twisted
tree, many trunks, forming
a thicket all by itself, on
low sand dunes.

2

28 Auricularia

on rotting Messerschmidia log
in mixed scrub forest.

3

29

on rotting Messerschmidia log
in mixed scrub forest

Sample #115 - polished
sand from ~~the beach~~
south passage beach at
about high tide level.

Bag. 120 - ^{just} seeds from Wotho
mostly from Eirek Islet, where,
for some reason, they were
infinitely more common than on other
islets.

8

March 19 - Omblim Islet
 sand platform, somewhat
 uneven surface, 1.5-2 m. ^{alt.}
 covered by a single large
 grove of *Pisonia*, medium
 sized, ^{perhaps 1.5 m. tall.} with undergrowth
 of root sprouts of *Pisonia* ^{alt.}
 This surrounded on all
 but southwest side by
 a narrow belt of *Messerschmidia*.
 Only other vascular plant
 is *Lepturus*, which forms
 a small patch on seaward
 side in ~~*Messerschmidia*~~ ^{*Tournefortia*}
 belt, now dry.

34430 *Pisonia grandis* R.Br.

Dominant tree, this one
 in *Messerschmidia* belt
 on lagoon side.

Peat layer well developed
 in central part ^{only} of *Pisonia* forest,
 from 1 to 3" thick, underlain
 by partially consolidated layer
 of sand 1-2" thick, this lying
 on loose sand. On top is a
 loose layer of dry leaves
 covered by guano stains.

Sample #116 - peat from a
 rather thick place.

Sample #117 - partially cemented
 sand from beneath #116.

Near center is a broad
 sand dune about 4 m.
 above high tide. Some
 sand is piled up around
 edges, burying lower parts
 of ~~*Messerschmidia*~~ ^{*Tournefortia*} trunks.
 Waves cutting a low cliff
 in sand on north side, exposing
 abundant ~~*Messerschmidia*~~ ^{*Tournefortia*} roots
 4-5 m. from plants. Birds are
 abundant and the dead leaves,
 branches, etc. are strongly stained by
 guano, but no noticeable
 accumulation of guano seen.

Shrub 2 m. tall (other trees
 up to 15 m.), completely
 defoliated (others in
 various stages of
 development of new
 leaves), loaded with fruit.

Sample 118 (2 bags) peat and
 partially cemented ~~to~~ sand
 from place where peat is
 thin, has hard black surface.

Sample 119 - piece of a slab
 of somewhat weathered
 cemented sand lying on
 surface in central part
 of *Pisonia* forest. Possibly
 represents an older, now broken
 up layer of same cemented
 material from under peat.

March 19 - Kabben Islet

- 34431 *Canavalia microcarpa* (D.C.) Piper
 7 abundant on lagoon side of coconut plantation that is much overgrown with brush and vines.

- 5 32 *blue green*
 on open sand in sparse scattered scrub at west end

March 19 - Orbelum Islet

- 5 33 *lichen*
 on bark near base of large *Pisonia* trunk

- 9 March 20 - ~~Kabben Islet~~ ^{Bigen Islet}
 sand platform with high beaches, general surface about 2 m. above high tide level, but uneven, one large dune hill being about 2 m. above general surface. Lagoon beach being cut away by waves, leaving a 2 m. sand cliff, with a pile of slabs of beach-rock at base. One slab up in center of islet on sand.

- 4 34 *Portulaca lutea* Sol. ?
 scattered tufts in coral sand under *Pisonia* trees.

extensive vine, tangled over shrubs and on ground; fruit immature.

forming a wrinkled crust.

- 9 Vegetation - rather open *Pisonia* forest in center, some fair sized trees, some bare, several still with last season's leaves. Occasional thin patches of *Lepturus*, scattered *Boerhaavia* and *Portulaca lutea*, latter nearly leafless, the thick stems carrying it over. Sparse belt of ~~*Messerschmidia*~~ ^{*Tournefortia*} around periphery, with a few leafy bushes. Many white-capped noddy nests in trees. almost leafless, old fruiting branches dead, remaining leaves purple; stem dull brown, fleshy.

7

March 19-20 - Kabben Islet

A large triangular island with one lagoon coast and two seaward ones, an east and a southwest. There are also two passage beaches, one north, the other west, the latter not sharply marked off from southwest seaward beach.

Most of the beaches are of sand or fine white gravel. Along the lagoon beach is a broad ridge of sand, broken into small dunes at the western end. The north passage beach is of pebbles, lying on beach-rock, changing to finer gravel seaward and around on seaward beach changes to sand. Pebble beach is backed by a low ridge of ^{beach} rock.

The south ~~point~~ peninsula of the islet is surrounded by a pebble-cobble beach, high and steep, backed by a very broad ridge of boulders and rubble, 100-200 m. wide. This beach rests on a pitted conglomerate platform with remnants of a higher platform here and there not yet completely eroded away on the southwest side. The

Vial #139 - animals collected in and under rotting Messerschmidia log in mixed forest, Kabben I., March 20.

7

ridge and boulder beach extend around the point and ~~for~~ to the beginning of the sparse scrub and the crescent of sand beach on the s.w. side and a corresponding distance, perhaps 300 m. or more, on the east side.

The southwest corner of the islet ~~has~~ has a broad reef-rock platform, and three series of beach-rock extend along the passage beach, dipping ~~down~~ away from the islet, splaying out fan-wise toward the lagoon corner, but not reaching it. This part has a sand beach.

Along the top of the lagoon beach is a narrow belt or fringe of scrub of *Sesuvola* with some *Messerschmidia*, tangled with *Wedelia*. This is in the edge of the coconut plantation, on the sand ridge.
(cont. on p. 90)

March 20 - Kabben Islet

34435 *Cordia subcordata* Lam.

4

large clump in mixed forest just in from lagoon beach - common in mixed forests around coasts of islet

36 *Turbinaria ornata* J. Ag. (det. Taylor 1955)
cast upon east seaward beach.

Sample #121 - drift seeds from Kabben I. Wotho Atoll (note of pentagonal *Barringtonia* fruit.)

Sample 122 - pumice from east beach, Kabben I. Wotho Atoll

Sample 123 - peat from under *Pisonia* trees, central part Kabben Islet. (from layer about 6" thick, loose below, many roots removed.)

Sample 124 - peat from same place as #123, but layer only 2-3" thick. Surface black.

Sample 125 - black crumbly soil with fine rubble from beneath #124. about 4-6" deep, on fine rubble.

large spreading tree; sterile.

dried fragments, bleached.

7

Under large *Pisonia* trees near center of eastern half of island, is a well developed layer of spongy peat. In places sampled it varies from 2 to 6 inches thick. Surface is black and smooth under a loose thin layer of dead leaves. No noticeable guano. Peat is very dry. Inside it is brown. Many large living roots are in the layer.

Under it is simply loose blackish crumbly soil with much small rubble. No noticeable sign of either consolidation cementing, or softening of rubble fragments.
(see samples 123-125)

Encoelma

March 21 - Moke Moke Islet

- 34437 *Caulerpa uvilliana* Mont. (det. Taylor 1955)
 well below low tide level
 on smooth rock bottom of
 passage west of islet
- 3 38 *Caulerpa uvilliana* Mont. (det. Taylor 1955)
 epiphytic on *Caulerpa* #34437
- 6 39 *Terminalia samoensis* Rech.
 in sparse mixed scrub-forest

March 21 - ~~south~~ north
 half of long ~~sand~~
 unnamed sand island
 on east reef

- 5 40 *Terminalia samoensis* Rech.
 occasional in sparse
 to closed scrub on
 sand dunes.
- 3 41 *Suriana maritima* L. ✓
 one very small colony
 on seaward side on
 low sand flat
- 7 42 *Pemphis acidula* Forst. ✓
 one small but very
 dense clump ~~on~~
 on whole island

(43 *Halimeda*
 bleached, washed up on beach

rhizomes creeping, straight

reddish

shrub 1 m. tall; flowers
 cream white.

shrub 2.5 m. tall,
 one almost leafless,
 but covered with
 red fleshy terete fruit,
 other, same clump,
 sterile but with leave
 seedlings on leeward
 side on soft sand.

shrub 1 m. tall,
 leaves bright green,
 flowers dried up.

densely branched
 shrubs 4 m. tall;
 thin leaves on lagoon
 side, thick fleshy
 ones on windward;
 flowers white.

March 21 -

Vial #137 - bottom layer - ~~insects~~ ^{animals} collected under loose bark of dead stub of *Messerschmidia* tree, north end of unnamed long sand spit on east reef.

second layer - small beetle, brushed off vegetation, Mokenok Islet, March 21. P-201431

Sample #126 - periwinkle scattered over surface of bare sand on Mokenok Islet. (2 bags) (see p. 77) P-201432

Vial #141 - Mokenok Islet - March 21, flies collected around large log cast up on beach, with masses of dead goose-neck barnacles which are possibly what attracted the flies.

Vial 138 - insects and spiders collected around large *Pemphix acidula* clump on long unnamed sand spit on east reef. Many insects seen on shaded side of clump (9-9:30 a.m.) none on sunny side.

V. 05
11

March 22 - Wotho Islet

East end surrounded by pebble or boulder beach, lagoon beach west of this, sand. Seaward beach, as seen from west, is a high steep boulder ridge.

Most of the islet is covered by dense *Pisonia* forest, reaching out to seaward edge. Surrounded on lagoon side, by a broken fringe

10

March 22 - Medyaron Islet

East passage beach, seen from east, very long. Outer third is boulder beach, middle third seems to be gravel, inner third sand. A well-developed belt of *Messerschmidia* extends along entire beach, backed by coconut plantation.

Lagoon beach is all sand. In the eastern part there is no very noticeable sand ridge. In the middle the sand ridge along the back

11a

~~*Tournefortia*~~
of ~~*Messerschmidia*~~.
A clump of coconuts at west end.

Seen only from boat in lagoon.

Sample #128 - drift seeds from Medyaron, Mokenols and sand spit on east reef.

10

of the beach is broad and low, but very definite, actually appearing to be two ridges. Westward, about the west third of it develops into conspicuous dunes from 6 to 10 feet high. Back of lagoon beach is a broad belt of open scrub, of *Scaevola*, ~~*Messerschmidia*~~, ~~*Tournefortia*~~, *Quettarda*, occasional *Pandanus* and *Buriana*. Between them is open sand. On this are

10

scattered pumice and scoria pebbles with network of roots on under side. (~~Hampl~~)

Here and there in this belt are great clumps of *Calophyllum*. Their leaves are browned on tips or edges where much exposed to weather.

On the dunes to the west the scrub is about the same composition as on the sand flats back of the beach to the east, but with much more *Luriana*. In openings *Triumfetta* is often found, half smothered by sand.

In other openings *Lepturus* is common and *Cassytha* is common both on *Lepturus* and on bushes. ~~to~~

To the west, near point the scrub becomes taller and denser, and may be termed scrub forest.

The island is extensively planted to coconuts, on a fine sandy gray-brown soil. The part of the plantation near the lagoon

10

beach ~~is~~ is rather well cleared out, with scattered small bushes, the ground covered by a mat of *Canavallis* or *Wedelia*. Large *Calophyllum* trees are found ~~at~~ around old house site and elsewhere where there might have been a house, just back from the beach scrub. (*Calophyllum* is more abundant on this islet than anywhere I have been in the Marshalls. Probably this accounts for the abundance of *Calophyllum* drift seeds on the beaches of various islets of Wotho).

Away from the lagoon the woody second story becomes more abundant, forming thickets, composed of *Pandanus*, *Morinda*, *Terminalia*, *Scaevola*, a little *Cordia*, tangled with *Ipomoea* ~~a~~ *tuba* and *Canavallis*, and choked

10 with coconut seedlings of all ages. Walking is difficult. There is some ground cover of *Polypodium* and *Peperomia* the latter especially in small openings. In thin places in the plantation ~~*Messerschmidia*~~ ^{*Tournefortia*} and *Guettarda* are very ~~abundant~~ common, with ~~*Wedelia*~~ ^{*Wedelia*} on the ground.

Natives visiting this islet, had gathered a bundle of dry Pandanus leaves and some copra and turtle eggs, as well as 3 or 4 dozen coconut crabs. Some of these had been roasted and the large claws tasted very good. A ~~one~~ large individual from this batch was preserved - in tank of alcoholic specimens otherwise collected by MacNeil. (unnumbered). The color varied on different individuals, dull purple usually predominating, but certain ones largely light blue.

12 March 21 - Mokermok (Mokermok) Islet - Belt toward lagoon a sand flat, forming 2 parallel ridges, especially toward the west, where there are two more ridges, separated from the two lagoonward ones by an interval, in which there is locally a little rubble. These ridges are generally sandy.

On the sand flat or ~~and~~ two lagoonward ridges is a sparse stand of scrub of *Scaevola*, *Guettarda*, and ~~*Messerschmidia*~~ ^{*Tournefortia*}. Much bare sand and much *Cassytha* in the bushes. *Purpurea* (sample #126) is scattered commonly on this sand. Some of the pieces have patches of a root network on the bottom.

The rest of the island is wooded with a mixed forest, in most parts rather sparse, made up of ~~*Messerschmidia*~~ ^{*Tournefortia*}, *Guettarda*, *Pisonia*, *Cordia*,

12

and *Scaevola*, the ~~*Messerschmidia*~~ ^{*Terminalia*} most abundant near the edges, the *Pisonia* almost dominant in center. Three sparse patches of coconuts are found near the lagoon side. They show evidence of the drought in having the recent clusters of nuts dwarfed or abortive, some nuts small on some of the older clusters. Trees in the westernmost clump mostly have no nuts. The forest is rather dense in center of widest part of island, where *Pisonia* is commonest, about 8-10 m. tall, little undergrowth or ground cover. To the west it thins out, becomes rather sparse, ~~*Messerschmidia*~~ ^{*Terminalia*} more abundant, *Pisonia* less, thin ground cover of *Lepturus* with some *Boerhaavia*, *Cassytha* especially in open places. One can walk freely almost anywhere. More or less same on

12

east end near passage beach. Some *Terminalia* here. The low rounded mounds of ^{broken} coral perhaps betray their origin here. A little to seaward of center of islet, where the soil is generally rather rubbly, low mounds of small pieces have piled up around root systems of *Pisonia* trees, either surrounding them or on the seaward side. In a few cases the tree has fallen, leaving the mound, with the tipped up root system adjacent, lying toward the lagoon either contiguous or several feet away. There are examples in various states of preservation, one or two having sent up big sprouts from the fallen trunk, one having almost decayed away, and several lacking any root system, the mound alone surviving.

12

Around the peninsula that sticks out to seaward, from near the lagoon end of the east passage beach around to the ^{interior} bend in the west passage beach, is a well developed boulder ridge, with *Caecula* - *Messerschmidia* scrub, ^{on the north east side,} this changing inward to the forest described above.

Back of the west end are several channels between rubble & reef conglomerate flats.

Water runs swiftly through these from seaward, where they are narrow. They are much wider toward the islet, then the current sweeps westward from each of them around the west horn of the islet.

In these the "oil streaks" described before were very prominent, following the direction of the wind rather than that of the current, though they are not very different. Opportunity was taken to

examine these carefully. No iridescence whatever was noted. The boundaries are extremely sharp. The phenomenon is one, certainly, of a much increased surface tension. There was a suggestion of difference in feel of the water, but not very tangible. No difference in taste. Attempt was made to follow several of these streaks to their sources, which seemed to be in the rubble flat to windward, but the appearance finally gradually disappeared as the source was approached. (MacNeil reports seeing small emissions of bubbles coming out of reefs in places). No origin obvious. Material did not seem to have come in from the sea, though this was not impossible here.

B. leucocaulis

3

Mar. 21 - Long unnamed sand spit on east reef. Examined north third of this.

The spit is a long low ridge of sand on platform that is generally marked with several series of beach rock, that showing along lagoon beach mostly dipping toward lagoon, that on seaward flat of reef conglomerate mostly dipping seaward. However, one stretch of a single series of beds, lying between the seaward dipping beds and the main sand ridge dips lagoonward, making it look as though a stretch of the sand ridge has moved into the lagoon. The denuded outer flats are generally rather wide.

There is very little vegetation, only a few scattered *Sonneratia* ~~*Sonneratia*~~ *Sonneratia* and *Scaevola* bushes, on the thin

3

sand locally found on these seaward flats. On the main sand ridge is a mixture of *Scaevola*, ~~*Sonneratia*~~ *Sonneratia* and *Terminalia*, varying in density from scattered to continuous, some ~~*Sonneratia*~~ *Sonneratia* trees 5-6 m. high and 2-3 dm. thick, but these uncommon.

Where there are scattered bushes low mounds of sand have accumulated ~~between~~ under them. Where there is continuous scrub the sand ridge is continuously several feet higher, with highest line usually nearest the lagoon side.

On seaward side is one tiny patch of *Sonneratia*, and about one third the way from the north end is a large and conspicuous clump of *Pemphis acidula*, quite dense. On the lagoon side the leaves

are rather thin, on the seaward side very thick. At 9-9:30 a.m. insects of various sorts were common on the west side of this clump (see ~~Bottle 138, p. 71~~) while none were seen on the sunny side. The *Pemphigus* was flowering.

The sand, where there is no vegetation tends to be flat on top. It is fine, white to pinkish, and the surface may be held firm by an incipient algal crust. There is a loose layer of dead leaves under the bushes. Here and there are thin patches of *Lepturus* in the bushy vegetation, none in open.

There is a small group of ragged coconut trees at north end, and a very few ragged individual trees here and there along the islet.

From the lagoon the vegetation all along seems to be scrub with frequent *Messerschmidia* 2-3 times as high as general level of scrub.

North of this spit are some small patches of sand on the reef. Then there is a small islet with two patches of vegetation connected by a rock flat. The vegetation of both halves is sparse scrub and there is one coconut tree on south half. (Seen only from boat).

Between this and the south end of Wotho Islet is a small bar on a rock platform. From a boat two bushes are visible on this.

2

March 18 - Wotho Islet -

The grass, and, especially, the ~~grass~~ *Fimbristylis*, show strongly the effect of the drought, by being brown and partly dried up.

Beach rock, dipping lagoonward, extends for some distance along south end of lagoon beach. It is pitted with rounded edges, probably abraded by wave-washed sand.

Toward the south end

3

perhaps 200-300 m. north of point, there is an abrupt change in the height of the coconut trees, those to the south being lower.

The south point has only mixed scrub and, back from point, mixed scrub forest.

March 18 - South of the long sand spit and to the lee of it, "oil streaks" were observed, not very numerous, but a few of them very distinct. Those to leeward of the sand spit would seem to preclude the idea that oil was coming over the reef from the open sea. Also common on both sides of Ertjeman!

Leg 14

Ruswaa Islet - seen only from boat in lagoon. On lagoon side appear to be high rock beds, possibly undercut by lagoon waves, extending above all but most extreme high tides.

Vegetation is a sparse scrub and scrub forest

On Ertjeman the lagoon beach is lined with great rocks with great slabs of it piled up on beach above. Some sand on north end, this with scrub of *Messerschmidia* and *Scaevola*.

4

mostly of *Messerschmidia* with some *Scaevola*, a few coconuts on lagoon side.

5

Ertjeman, Uditj, and Jabenau Islets - seen only from boat in lagoon.

These are all wooded possibly with *Pisonia*, as defoliation is very noticeable, but from appearance on aerial photos, rather mixed. The

ends and in case of *Ertjeman*, the seaward projection tapers off to scrub. On *Jabenau* MacNeil reports a single *Pemphis* bush on extreme end of seaward projection, the rest of the scrub being *Messerschmidia* and *Scaevola*. This projection is bare rock.

On each of these islets there is a small patch of coconuts on the lagoon side.

Lagoon side of Ertjeman mixed forest of *Pisonia*, *Messerschmidia*, *Guettarda*.

4 March 19 - Eniwetok?

The entire lagoon beach, except for a few short stretches, is lined by lagoonward dipping beach-rock, and above this, great slabs of beach rock are piled up to the top of the beach. The unusually strong lagoon waves are dissolving the beach-rock along bedding planes, cracks, etc. the cracks apparently result of undercutting along bedding planes.

A slight ridge is developed here and there along lagoon beach, but it is not conspicuous.

Along the entire lagoon beach, except for a short stretch of coconuts with *Scaevola* between them, is a mixed forest of *Guettarda*, ~~*Messerschmidia*~~ ^{*Tournefortia*}, *Pandanus*, *Pisonia*, ~~*Occhrosia*~~ ^{*Messerschmidia*} and *Scaevola*, tangled with ~~*Wedelia*~~ ^{*Wedelia*} and *Ipomoea pes-caprae*. Here and there are small patches of *Lepturus*.

Inward *Pisonia* becomes dominant, but is being

6 replaced by a vigorous layer of young ~~*Occhrosia*~~ ^{*Messerschmidia*} trees.

In the center, at least of the south end, is a solid forest of ~~*Occhrosia*~~ ^{*Messerschmidia*}.

In all of this forest sticks blackened by *Xylaria* are common. *Rat* are common.

The soil in the forest is fine and blackish, ~~gotten~~ with varying ~~quantities~~ admixture of fine rubble.

Toward the seaward beach it becomes more sandy and has on the surface boulders strewn into the forest.

~~At~~ the south end is mixed forest similar to that described for the lagoon beach, with, also, some *Canavalia* tangled in it. This peters out into mixed scrub on the sandy extreme south end.

The entire seaward beach is gravel, resting on flat reef conglomerate. Above it is a belt of ~~*Messerschmidia*~~ ^{*Tournefortia*} forest with a *Scaevola* fringe.

72.

Kabben Islet (ctd. from p. 105)
The plantation, except immediately around the old hut, is choked with thickets. Near the lagoon these thickets are of Pandanus, Scaevola and Morinda, tangled with Wedelia. Around the hut, where the thickets have been cleared, a thin tangled mat of Wedelia covers the ground.

West of the center of the lagoon beach, ~~Delonix~~ becomes a component of the plantation thickets, ^{very} locally forming almost pure stands. From here westward the coconuts are very sparse, thickets dense.

Back from the beach, in the region of the hut, Scaevola rapidly drops out. Quettarda, Morinda and Pandanus dominate the thickets and the whole becomes choked with young coconuts of all ages.

Still further back the coconuts become sparse. There are large clumps of breadfruit, and ~~Occhrosia~~ ^{Macaranga} becomes common.

with
quettarda
and Pandanus
seedlings
making
under
growth.

7

Westward the plantation gradually changes to mixed scrub forest with a few coconuts. On the west end, behind the low dunes, there is open sand with a scattered scrub of Scaevola, Quettarda, Luriana, and ~~Messerschmidia~~ ^{Terminalia}, with scattered trees of Pandanus and small coconut trees. On the sand is a well-developed algal crust.

On the north passage a belt of scrub, mostly Scaevola and very low on the seaward end, becoming mixed with ~~Messerschmidia~~ ^{Terminalia} and Quettarda, and taller, toward lagoon, occupies the ridge of broken coral. Back of this is a scrub forest of Messerschmidia, Pandanus, Pisonia, Quettarda, Soulaumea, with a little Scaevola and Terminalia. Locally there is a ground cover of rather wilted Polypodium, and of Soulaumea seedlings, these now mostly dead. A few saplings of the trees and a tangle of ~~Wedelia~~ ^{Wedelia}

7 forest extending inward from the east coast is reached. The soil remains the same uneven mixture of loam and fine rubble. In the mixed forest around the pure *Ochrosia* ~~*Wickstroemia*~~ behaves as a liana, straight stems going up into the canopy, then forming tangles.

Extending from the east coast in for some distance is a great ^{roughly} triangular area of pure ~~*Ochrosia*~~ ^{*Messerschmidia*} forest. This has a ~~poor~~ ^{thin} layer of fruits on the ground and a continuous layer of seedling, varying locally in height from 2 dm. to 1 m. The height of the forest varies from perhaps 20 m. ~~down~~ sloping down to the east to perhaps 4-5 m. there it gives way to a narrow belt of scrub, mainly *Caeculia*, which lines the beach. There is some admixture of *Guettarda*, and locally clumps of ~~*Messerschmidia*~~ ^{*Tournefortia*} trees. A characteristic of this scrub belt on the windward beach is the effect

7 of strong grooves or channels in the surface of the vegetation, parallel to the wind direction, starting at the beach and becoming shallower and disappearing inward. This scrub belt extends the full length of the east coast, typically wedge shaped in cross section, thin edge to the beach.

Along the inner sides of the ~~*Ochrosia*~~ ^{*Messerschmidia*} forest great *Pisonia* trees are mixed with the ~~*Ochrosia*~~ ^{*Messerschmidia*}. Here, also, are many patches where the *Ochrosia* is very chlorotic, and even some trees have died.

Just inland from the ~~*Ochrosia*~~ ^{*Messerschmidia*} forest is a mixed forest of great *Pisonia* trees, a *Ochrosia* and a few old ~~*Messerschmidia*~~ ^{*Tournefortia*} trees, a few *Pandanus* and small *Allophyllus*. Undergrowth is of saplings of ~~*Ochrosia*~~ ^{*Messerschmidia*} and a few of *Allophyllus*. There are occasional scattered

7

groups of the low rounded mounds of broken coral previously described. One old ~~Messerschmidia~~ ^{newspine} had fallen, laced off. it was about 45 m. tall. A few of the ~~Achrosia~~ ^{newspine} here are old, but there is mainly a stratum of young ones about 10 m. tall. The undergrowth is of young ~~Achrosia~~ ^{newspine} saplings, and a few young Pandanus.

Northward near center of this part of the island is a sparse stand of coconuts choked with Guettarda, ~~Achrosia~~ ^{newspine}, Pandanus etc. with seedlings of these and of coconuts making walking difficult.

Then there is a section of scrub forest of Guettarda, Bougainvillea, Pisonia, and Pandanus.

Between this and the east end of the lagoon beach is more choked coconut plantation.

7

The east coast scrub belt sends a narrow hook around the south point. This is backed by mixed forest, the transition being gradual. Guettarda and ~~Messerschmidia~~ ^{newspine} replacing more and more of the ~~Acrova~~ ^{newspine}.

Along the boulder beach of the southwest coast of the south peninsula mixed forest comes right to the top of the beach. It is made up of Pisonia, ~~Messerschmidia~~ ^{newspine}, Guettarda, and Cordia.

Inland on the broad boulder flat Pisonia becomes dominant in this mixed forest, some of the trees being enormous. Inland from there is a large opening on the rubble flat filled with a deep mat of ~~Wedelia~~ ^{newspine}. In the inner edge of this is a grove of coconuts.

The boulder and rubble flat along this coast is 100-200 m. wide.

Along the central part

7

of the southwest side is a wide sand flat with very sparse scrub, much open sand, *Scaevola* predominant.

In the mixed forest on the rubble and boulder flat there is little undergrowth except occasional seedlings of the component trees and, inland, of ~~*Acacia*~~ ^{*mesquite*} which is rare as a large tree but is actively invading.

The *Cordia*s, abundant near the beach, are enormous low spreading trees with immense gnarled trunks and great withering horizontal branches just off the ground. Penetration difficult.

8

March 20 - Enebek Islet

The surface of this islet is all sand. Along the lagoon beach is a broad dune ridge from 2 to 4 m. above high tide level. The seaward part is lower.

On top of a high rounded dune on the south end ~~is~~ are scattered large boulders, about 2.5 - 3 m. above high tide.

On this dune there is a grove of *Pisonia*, surrounded by ~~*Messerschmidia*~~ ^{*Terminalia*} and *Terminalia*.

Both the *Pisonia* and the *Terminalia* were practically leafless. North of this a large *Cordia* forms a broad low tangled thicket.

In the center are three patches of coconuts, the middle one back from the beach. Between them is a patch of ~~*Acacia*~~ ^{*mesquite*}. Here there are, for a short distance, two dune ridges, but they are lower than at the two ends. The west ~~north~~ end and parts of

1952 Marshall Is.

the ~~south~~ east and center are covered by a mixed forest of ~~Messerschmidia~~ ^{*Sonneratia*}, *Pisonia*, *Terminalia*, and *Guettarda*. The edges of this and sparser spots are tangled with ~~*Wedelia*~~ ^{*Wallingtonia*}. Some of the *Terminalia* has lost its leaves, some has not.

The lower ground back of the dune ridge is covered by uneven scrub forest, mainly *Messerschmidia*, *Guettarda*, and *Pisonia*, of ~~var~~ varying density.

Toward the seaward beach, especially on the ~~east~~ ^{east} and ~~west~~ ^{west} ends, this thins out to a sparse scrub of *Scaevola* with some ~~*Messerschmidia*~~ ^{*Sonneratia*} and some *Guettarda*. Much bare sand between the rounded bushes.

A broad series of beach-rock, dipping seaward, lines the southeast beach. Some beachrock also, on the south side, and along the southwest side.

Wotho Atoll

Drift seeds are unusually abundant on this island.

The sand at about high tide level, and the beach sandstone beneath it, are curiously polished. (sample 915, p. 58).

13

Wotho Birds:-

In general, birds are abundant on the smaller islets, relatively scarce on the larger ones. Nesting is mostly seen on the smaller islets. Terns, as usual, are the most abundant birds.

Frigate birds -

a colony of well over a hundred roost on Ambelim Islet. During the hot part of the day they are mainly resting, toward evening they soar in numbers above this and other islets, or out over the sea. Several individuals are all black beneath. Some also seen over Bigen Islet.

Terns - Both noddies, fairy terns, and rarely black capped black-naped terns may be found fishing together, flying excitedly over schools of fish, diving for them, either in lagoon or out in open sea. After they are full the noddies rest in the water.

13

Brown booby -

One seen several times around Bigen Islet. An immature one seen at sea west of Bigen. One adult seen at Ambelim Islet.

Common noddy -

Seen in small or moderate numbers over lagoon and at sea, a few on the larger islets. Many on Eriherudo Islet. Many birds and several nests seen on Eirik Islet, with downy young. One bird scared up here that had difficulty flying because it had *Pisonia* fruits stuck over its feathers. Several nests on Bigen Islet. Many birds seen on the long sand spit on the east reef.

Some birds and one nest on Mokemok, this low in a *Guetarda* tree.

Commonly seen, in middle of day, in groups on beaches.

13

White-capped noddy -

Commonly seen fishing over lagoon or open sea, at any time of day. Some nests seen in *Schinus* trees on Enikerudo I., also about 20 nests on Eirek. Hundreds of nests and a multitude of birds on *Ombelium* in Pisonia and also in *Messerschmidia*, many seen holding a "conference" in *Messerschmidia* trees on seaward side. On the seaward side of Kabben Islet were a small group of nests in Pisonia trees. Many nests in Pisonia and *Messerschmidia* on Bigen I.

Fairy terns -

To be seen generally almost anywhere, flying over lagoon or open sea in pairs or small groups, or fishing in flocks in company with other terns, or flying overhead on islets protesting intrusion into their privacy. Probably in most such cases there are eggs on

13

the bare branches of the trees, but they are seen only with difficulty. Many seen on *Ombelium* Islet, ~~Enikerudo Islet~~ ^{on the latter} where one egg was seen on a *Messerschmidia* branch. On Enikerudo many were flying overhead, protesting as though they had eggs in the trees. Many were flying over the long sand-spit on the east reef, also over Moke-moh Islet.

13
saw one
half
feathered
and one
fully
feathered
young.

Black-naped tern -

Small flocks on various islets, invariably on or over sand projections or beaches on inner corners of islets. On Eirek there were 3 on the east corner and 4 on the west. On *Ombelium* there were 10 birds on the inner beach - the island is too short for the corners to be distinguished. 6 were seen fishing over the lagoon near Wotho at 6:30 p.m. and

one with a flock of other terns near Moke-moh at about 4 p.m.

13

Crested tern -

Rare here, possibly only one or two pairs.

One seen at sea west of atoll. Two seen on Imbelin Islet.

Pacific Golden Plover.

Seen in small numbers generally, on inner and outer beaches, and in interior wherever conditions are at all open, on most islets. Often with turnstones, tattlers, or curlews.

13

Wandering tattler -

Seen one or two at a time, usually on outer beach or reef flats, or passage beaches on most islets. Often in company of plovers or turnstones, or more rarely, curlews.

Turnstone -

In small flocks of two to five on most islets, on beaches or reef flats. Often with plovers and tattlers.

13

Bristle-thighed curlew -
 seen singly, in pairs,
 or as many as three or four,
 on beaches and reef flats
 of most islets. Here
 often fairly tame,
 allowing one to approach
 to within 15 or 20 m.
 before flying.

Whimbrel -

Rare. One seen on ^{Eniwetok} Eniwetok Islet.

Itinerary of voyage -
from Capt. M.W. Lhoaf's log book
of 75367.

Left Yokohama Nov. 16 - 13:51

arr. Taongi atoll " 25 - 06:30

Dep. Taongi atoll " 25 - 16:00

arr. Bikar " " 26 09:30

Dep. " " " 26 11:30

arr. Utirik " " 26 17:30

Dep. Utirik " ~~Dec~~ 28 12:30

arr. Kwaj " " 29 10:00

Dep. Kwaj " Dec. 1 16:00

arr. Utirik " " 2 15:00

Dep. Utirik " " 4 14:00

arr. Taka " " 4 15:30

Dep. Taka " " 9 12:00

arr. Utirik " " 9 13:15

Dep. Utirik " " 10 10:30

arr. Ailuk " " 10 15:30

Dep. Ailuk " " 11 08:15

arr. Likiep " " 11 13:15

Dep. Likiep " " 18 09:00

arr. Jemo " " 18 11:00

Dep. Jemo " " 18 14:00

arr. Likiep " " 18 17:00

Dep. Likiep " " 23 08:30

arr. Jemo " " 23 12:30

Dep. Jemo " " 23 15:00

arr. Ailuk " " 23 18:00

Dep. Ailuk " Jan. 4 08:45

arr. Lae " " 5 09:30

Dep. Lae Atoll Jan. 5 20:30

arr. Kwaj " " 6 08:15

Dep. Kwaj " Feb. 1 16:00

arr. Ujelang " " 3 10:00

Dep. Ujelang " " 10 16:30

arr. Wotho " " 12 09:00

Dep. Wotho " " 16 02:30

arr. Ujae " " 16 09:15

Dep. Ujae " " 24 21:30

arr. Kwaj " " 25 10:00

Dep. Kwaj " Mar. 1 07:30

arr. Ujae " " 1 18:30

Dep. Ujae " " 14 20:00

arr. Kwaj " " 15 09:30

Dep. Kwaj " " 17 19:00

arr. Wotho " " 18 10:00

Dep. Wotho " " 23 07:30

arr. Bikar " " 24 18:30

Dep. Bikar " " 24 18:30

arr. Taongi " " 25 9:00

Dep. Taongi " " 25 20:00

arr. Kwaj " " 27 10:30

Dep. Kwaj " Mar. 31 3:15

arr. Guam I. " " 10:30

Dep. Guam I. Apr. 2 8:00

arr. Tokyo " " 2 5:00

March 24 - Bikar atoll
from ship at from $\frac{1}{4}$ to $\frac{1}{2}$ mile.

Passage at 10:30^{am} - low tide,
quiet water outside, entire
reef showing, edges of
passage out of water,
conspicuously undercut.
Strong current flowing
out + 2000 rpm of penyan
boat ~~go~~ engine just held
boat in place.

at 3:30 pm. half tide -
no noticeable current,
but small swells
across mouth of
passage cause tide-rip
and considerable turbulence,
suggesting some current.
Wind waves from inside
may have caused some
of the turbulence.

at 6:30 pm. - high tide -
or going down somewhat -
tremendous current
coming out, making
a wake a couple of
hundred yards out to
sea. Great swells
breaking across. Between
swells the channel like
a mill race.

Leeward reef has
numerous boulders
strewn over it.
Windward reef, north
of islets, has a few,
but larger.
North part of reef
flat very wide,
a sand bar on inner
edge.

~~Tall~~
Jaboverukku Islet -
covered by tall *Pisonia*
forest, trees partly
defoliated. A second
layer $\frac{2}{3}$ - $\frac{3}{4}$ height of
canopy, much greener,
unidentified, slopes off
to windward into scrub.
May also be *Pisonia*. A
sparse fringe of *Messerschmidia* along both
inner and outer beaches.
One greener clump possibly
Quettarda.

~~Tall~~
Between Jaboverukku and
Armenia islets a
nearly continuous
sand and rubble flat
a rock platform. On
highest parts of this a
very few young *Messerschmidia*.

^{Alum}
Arumenis Islet - on northern projection a sparse stand of *Messerschmidia*. Main part of islet a *Pisonia* forest with sparse fringe of *Messerschmidia* around edges. Some small sand dunes on south side.

Jaboera Islet - an unvegetated sand bar held by beach rock on seaward side.

Bikar Islet - mostly tall *Pisonia* forest, with a belt of very sparse scrub around it especially on north and south ends. Scrub mostly *Messerschmidia*, a very few *Scaevola* bushes. Several coconut trees in outer part, a large grove ~~of~~ of them on inner end, between forest and scrub belt. Dunes on south side. On south side some enormous boulders on beach.

"Oil streaks" numerous to lee of south half of atoll - rather braided and broken up at bet. $\frac{1}{4}$ & $\frac{1}{2}$ mile, but some still fairly strong.

Birds -

Very few birds seen on leeward side of atoll - only a few brown boobies.

On north, east, and south, larger numbers of boobies, brown, red-footed, and a few blue-faced, fishing with some noddies (common?) and a few fairy terns.

Great numbers of boobies, also many frigate birds roosting in trees on islets - scared up in great swarms by whistle blasts from ship at half mile.

Two unidentified fairly large gray terns seen flying.

The backs of brown boobies are sooty-brown, rather than chocolate brown as described by Mayr. Immatures rather gray-brown on backs. Three of these made passes at flag fluttering from masthead. Also dives for fish lure trolling behind ship.

March 25 - cruised around ~~a~~ atoll, sometimes within $\frac{1}{4}$ mile.

Vial 140 - parasitic insect taken off of *Lula leucogaster* caught on fish line $\frac{1}{4}$ to $\frac{1}{2}$ mile off Kamome Islet.

Near the northern end of the leeward reef, mostly opposite a reentrant angle in the reef, ~~is~~ a group of great boulders on the reef, 2 m. or more high, one, at least, several times as long, tending to be mushroom shaped because of a prominent top. ~~Some~~ some are tipped over. On windward side many small ones.

"oil streaks" such as described for Ujae, Wotho, Bikar, are conspicuous on leeward side of leeward reef, especially in the neighborhood of the passage. Extend for several miles, becoming very diffused.

All of the vegetated islets have boulder & cobble seaward beaches, material becoming finer southwestward. Large boulders strewn on reef and beach and up on islands.

Three northernmost islets have ~~very~~ sparse grayish scrub of *Messerschmidia*, innumerable birds. Vegetation on seaward side of Bibyll Islet not so sparse, mainly of low *Messerschmidia*, some scattered *Scaevola*. ~~On Pohak Islet~~ some areas where passages have filled, or perhaps vegetation has been swept off by storms, colonization is taking place.

On Pohak Islet there is more *Scaevola*. Also some taller trees here, probably all *Messerschmidia*, but possibly a few *Pisonia*.

South Islet absolutely bare of vegetation. Mostly cobbles locally gravel, great boulders strewn about.

Some
Messerschmidia
very
dry
looking,
dead or
appearing
so.

Pohak

The passage, at low tide, much like a mill-race. Water rushing out, current showing well out from mouth.

At almost high tide not much current. However, west swell had set in and breakers stretched across mouth. On day without swell this would be time to enter. Even with swell, periods of quiet observed as much as 10 minutes.

In neighborhood of mouth of passage a peculiar lighter blue color, suggesting effect of colloidal CaCO_3 in marl pools. Possibly fine CaCO_3 sediment coming out of lagoon.

no white seen on head above beak.

Birds -

One black-footed albatross seen ~~by~~ by Ch. Frey, on arrival in morning.

A large shearwater observed in flock at north point, sitting on water, and in numbers at various points flying. May be white-faced shearwater, but identity not certain. Dark sooty-brown above, white beneath but with wing primaries and secondaries, front edge of wing, and tail black. Irides with black smudged somewhat beneath also.

A smaller, but still large, black petrel or shearwater seen, not identified, flying close to waves on windward side in company with the other one.

Up to 8 or 10 frigate birds at a time following ship.

All three boobies present. The brown in large numbers. The immature ones, especially, following the ship and diving at trolling lures, several getting caught. (collected parasites off of one. They bite vigorously.) Blue faced and red footed in smaller numbers observed flying or fishing.

Sooty terns common flying around, sometimes fishing.

A few fairy terns flying.

Two white reef herons flew out from ^{leeward} reef, at different points, to ship, then flew back. One of these from near northern tip of atoll.

Fairy terns, sooty terns brown boobies, blue faced boobies, and frigate birds, seen fishing together.

plants seen but not collected.

Terminalia catappa

Pedilanthus tithymaloides

Gossipium

Bougainvillea ?

Pithecellobium dulce ?

Coccolobis uvifera

Pseuderanthemum atropurpureum

Cremum pedunculatum

March 31 - Guam -
much of forest on plateau at north end looks very decadent - large trees dying back. Forests on terraces below cliff-tops on east side, n.e. of Mt. Santa Rosa, and on north side, at Tulaque, are green and healthy-looking.

Around Mt. Santa Rosa are areas, just inside contact of volcanics with limestone, are wet depressions with standing water in rainy season, and places where water disappears under the limestone.

In quarry just north east or east of Anderson field pieces of limestone showing numerous Halimeda fragments were found.

Toward sea from this quarry is a very pronounced rim or wall at top of cliff, perhaps 6-8 m high. A hole has been blasted through this, giving an excellent view of the terraces below. On the face

of the cliff, just north of the hole, is a low grayish dense bush that I could not identify from distance, even with binoculars.

From air, it was possible to see dense growth of coconut seedlings around coconut trees in some areas on northern part of island.

April 1 - Guam -

Road down to Apra Harbor from Com-Marianas - Casuarina quite luxuriant, especially on road-cut, and less so away on sword-grass slopes. Some of these rather densely covered.

Road across from Base 18 to Talofofo - scattered Casuarina on most sword-grass areas. Some recently burned over north of road.

Fena River Dam area - Brookhart says most of forested area in valley bottom and lower slopes, as well as the belt across upper slopes of Fena drainage are on limestone

a debris strongly contaminated by limestone. Good stands of *Cananga odorata* observed on limestone slopes.

Extensive burning observed in Merizo area. Where not burned, *Casuarina* common on sword-grass slopes.

Rocks denuded by 1949 typhoon, south of Inarajan, fairly well covered by low *Pemphis*. Breadfruit trees that were, in Jan. 1950, denuded of leaves and small branches, now look almost normal again.

Coconut groves east of Merizo show very few effects of typhoon, and mangrove swamps show none at all, except lack of large trees. Not examined very carefully. In coconut groves there is now considerable undergrowth and well grown banana planting.

In general, driving around coast road, if one did not know there had been a typhoon and did not look especially for evidences,

one might not realize there had been one.

Padina is common on bottom at moderate depths of south end - up to 10 m. Considerable other algal growth at such depths. No *Halimeda* seen.

Large "vegetation" of soft corals in 1-2 m. of water in lagoon at end.

Kodachromes, taken by Brookhart, of Uracas, just after eruption in early 1952, show no vegetation on the cone or the lava flows. On the two old remnants of a former caldera that still exist at the edge there is green vegetation - possibly grass or *Peperomia*. Brookhart confirms that there was not a bit of green visible anywhere else on island, and that this vegetation was of very low stature.

April 2 - Around air field -

Weedy ground of a yellowish brown ash or tuff immediately surrounding air base and field. On rough ground and that not affected by traffic is a dense scrub of *Leucaena glauca* with occasional bushes of *Trema orientalis* v. *argentea*.

This scrub is from one to three or four m. tall. Occasional plants of *Acacia confusa* int.

Isthmus of black ash, mostly grassland with bunches of a large grass - *Miscanthus*?; ~~and~~ scrub in rough places, bright green so not that noted above. ^{Some bare ash, or with sparse *Canavalia*.}

Lunabachi has patches of scrub, probably *Leucaena*. patches of bare ash

- 34444 *Emilia javanica* (Burm.) Rob.
occasional around air base
45 *Phyllanthus urinaria*
rare around air base

Weeds observed immediately around air base:

- * *Cynodon dactylon*
- * *Tridax procumbens*
- * *Leucaena glauca*
- * *Sida rhombifolia*
- * *Crotalaria mucronata*
- * *Solanum nigrum*
- * *Pisum communis*
- * *Euphorbia hirta*
- * ~~*Phyllanthus*~~
- * *Vernonia cinerea*
- * *Chloris inflata*
- * *Lonehus* sp.
- * *Cyperus javanicus*
- * *Digitaria pruriens*?
- * *Euphorbia thymifolia*
- * *Oxalis corniculata*?
- * *Euphorbia heterophylla*
- * *Emilia javanica*
- * *Commelina* sp.
- * *Canavalia* sp.
- * *Achyranthes aspera*?
- * *Cassia leschenaultiana*
- * *Ticholaena repens*
- * *Agave sisalana*
- * *Phyllanthus urinaria*
- * *Cenchrus spinata*

flowers bright red.

> + * abundant
unrecorded from Two
firms. T. T. T. T. T.

April 13 - Road from Kumagawa to Chichibu - On the flat plain west of Kumagawa are prosperous farms, raising vegetables, millet, wheat, and especially mulberries for silk culture. Rice is doubtless produced in quantity in season but is not evident at this time.

The method of culture of the mulberries is a stump sprout culture, ~~with~~ with stumps a foot or two high, producing whip-like sprouts 1-2 yards long and very large leaves.

As one enters the mountains they rise rather abruptly and are covered by a low deciduous forest, a few m. tall, of mixed composition, apparently harvested by a coppicing system, as the trees are really clumps of ~~stump~~ stump sprouts. Scattered over this are widely separated evergreens and small clumps of evergreens. These become more prominent as one goes toward Chichibu.

There are many cherry trees as far as Nagatoro, and they are at their best today. In the lowlands below they have already lost most of their petals.

On bluffs and steep slopes are splashes of bright rose-purple, from masses of azaleas. It is a deciduous form with a wide open tube. These are especially prominent on a sharp ridge between Nagatoro and Chichibu where I took some photos.

April 13 - Road from Chichibu to

This has, basically, the same sort of vegetation as on the other road, but evergreens are more abundant, and in larger patches. A species of *Clethra* (?) with pendent inflorescences is prominent in the coppice.

April 19 - talked to Mr. Schultz, post-central man for CAA who has been here for ^{over} a couple of years and with Ruby Cox, nurse, also here for 15 months or so. The latter has some house plants in pots:

Scindapsus aureus
Cordyline terminalis
Dieffenbachia
 growing outside, close to house.

Catharanthus roseus
 (luxuriant where sheltered)

Criminum

plants small, densely caespitose, very purple, not flowering, somewhat sheltered.

Ananas comosus.

crown growing in box of soil, somewhat sheltered. *Polyscias guilfoylei* reached a couple of feet in calm season, then was cut back almost to ground by wind. Sweet potatoes grew luxuriantly where sheltered, any part exposed to wind was cut off instantly. Bugs ate the whole lot. *Casuarina equisetifolia* two saplings planted

behind wind breakers - one wind break fell down and that tree was almost killed. This grew a half meter in 6-8 months.

Three coconuts planted at same time, with different degrees of shelter. The more shaded one watered most. The most exposed is 2-3 dm. tall. The second about 1 m., the third 2 m.

Tomatoes said to do reasonably well.

Melons grow, but not well, fruit very small.

Three sp. of mosquitoes

Culex

Aedes aegypti

Aedes scutellaris or *albopictus* (?)

were common, now rare because breeding places destroyed.

5 red-bellied lava sparrows or pinches in captivity

Gambusia, introduced 4 years ago into a concrete reservoir, now abundant in brackish pools.

Flies now scarce, formerly abundant.

4 sp. rats formerly very abundant, millions, now not common.

not clear

April 20 - Wake Islet

Vial 141 - ants (*Solenopsis*)
and cochineal scale, on
~~roots of~~ crowns of
Setaria verticillata.

Vial 142 - dermestids from
a dead and dried tern
on ground; lice from a young
tern in pinfeather stage.
Black ant (~~from plant~~)
on floor in hotel.

Vial 143 - Odonate larva,
and *Gambusia* (mosquito
fish) from brackish water
in bomb crater.

Vial 144 - skink (lizard)
caught on ground in open
place, fairly common.

Vial 145 - flies caught
around garbage dump,
grasshoppers in open
grassy places, spiders
around buildings.
larva in grassy spot.

Mr. Frederick Schulty
3919 Waialae Ave.
Honolulu 55, T.H.
(ph. 77086)
(present c/o CAA Wake I.)

formerly ^{chief} Sanitation
Division, Territorial Board of
Health, T.H.

Now, temporarily, running
pest control work in
Wake I. for CAA.

Started - Feb. 14, 1951.

took 1 month to get started.

poisoned rats

emptied out mosquito

breeding places.

cleaned up garbage pits

April 20 - The entire atoll has been altered from the picture given by Bryan, who made his observations in the summer of 1923. Only in about three spots are there any noticeable number of trees that ~~seem~~ seem to date from pre-1941 time.

One of these is on the north leg of Wake Islet, west of the housing area. Here are large *Pisonia* trees, with some admixture of *Cordia*, little undergrowth. The trees show some "windshear" effect, and are only 3-10 m. tall. Trunks up to ~~several feet diam.~~ ^{8 dm.} diameter ~~at~~ ^{near} base but mostly much smaller.

The soil is of coral pebbles but locally the top layers show an accumulation of black peaty material, which, however, is quite friable. This area was relatively little torn up by military activities.

On the eastern side of the lagoon, between the housing area and the Pan Am compound are some areas of *Messerschmidia* that look as though they

are remnants of the pre-war forest.

The remaining such area is on the south ~~west~~ ^{east} leg of the island, just east of the Trans Ocean compound, where there is a small area of large *Messerschmidia* trees which stand out when observed from the air. Just east and lagoonward from this is some large *Pemphis* that was probably a part of the ~~pre~~ original stand mentioned by Bryan.

Around the northeast corner of the lagoon are still the saline flats of fine sand, mentioned by Bryan, with *Sesuvium* mats and especially inward, low tangles of *Pemphis*. This *Pemphis* is rather small and looks young, but ~~is~~ possibly remains from the stand described by Bryan. If so, it has been altered somewhat, opened up in places, grown up much taller, in places, than what Bryan reports.
(see p. 142)

Islet
April 20-21 - Wake Island
coral gravel and sandy soil

34446 *Amaranthus gracilis* ^{viridis} L.
around abandoned Japanese garden

2 47 *Pluchea odorata*
dominant plant forming
scrub in denuded or open places

3 48 *Nicotiana tabacum* L.
rare around abandoned
Japanese garden

3 49 *Ipomoea batatas* L.
rare around abandoned
Japanese garden

2 50 *Cynodon*?
~~not~~ common locally around
old Japanese garden, mostly
dead at this time of year

9 51 *Abutilon Indicum* (L.) Sweet
one plant near old Japanese
garden.

2 52 *Amaranthus tricolor* ^{dubius Mark.}
rare around old Japanese garden

9 53 *Boerhaavia diffusa* L.
common generally, especially
in open places (with 34454)

3 54 *Boerhaavia*
common generally, especially
in open places (with 34453)

1 55 *Euphorbia prostrata* Ait.
rare in open gravel

2 56 *Catharanthus roseus* (L.) Don
rare in open places

spreading herb

shrub 2 m. tall,
fls. pale lavender.

herb 1.5 m. tall; limb
of corolla bright pink.

shrub 1 m. tall,
flowers orange, faded
by 10 a.m.
- not spiny

prostrate, mat forming,
stems red, leaves dull
green above, white beneath,
flowers pink.

- prostrate, mat forming,
stems green, leaves green
above, white beneath,
flowers white.

prostrate, purplish.

flowers magenta.

- 34457 *Fimbristylis cymosa* R. & S.
2 abundant on recently
graded air strip border
- 1 58 *Eragrostis poaeoides* Beauv. ex R. & S.
rare along edge of air strip
- 1 59 *Eragrostis amabilis* (L.) W. & A.
along air strip
- 1 60 *Portulaca sarraensis* v. Paell.
rare in open place.
- 3 61 *Lesquerium portulacastrum* L.
common generally on low
saline places
- 3 62 *Lepturus repens* (Forst.) R. & S.
common locally in open
places. (Wilkes 924)
- 1 63 *Cyniza bonariensis* (L.) Griseb.
rare along roadside. (Wilkes 9.)
- 1 64 *Lonchus oleraceus* L.
rare along roadside (Wilkes 9.)
- 2 65 *Euphorbia hirta* L.
common around old buildings
- 1 66 *Ficus carica* L.
one plant around building
- 3 67 *Ficus rubiginosa* Desf.
one plant ~~in~~ around building
- 7 68 *Portulaca lutea* Sol.
very common generally
- 5 69 *Gossypium hirsutum* var. *religiosum* (L.) Watt
occasional, especially
around old Japanese installations.

mostly dead.

prostrate, fleshy.

prostrate, fleshy,
leaves somewhat compressed,
~~stems~~, glossy; stems
red; flowers rose pink.

flowers yellow

shrub 2 m. tall, fruiting
abundantly.
shrub 2 m. tall, sterile.

stems gray, ascending,
plant green, flowers
1.5-2 cm broad, yellow. stamens
25-30

spreading shrub with
elongate branches; flowers
cream-yellow.

- 34470 *Setaria verticillata* (L.) Beauv.
 1 very local, one patch in second growth.
- 2 71 *Coccolobis uvifera* L.
 rare in second growth around old installation.
- 6 72 *Pisonia grandis* R.Br.
 common generally, dominant locally
- 3 73 *Sporobolus per-caprae* (L.) Roth
 patches locally
- 1 74 *Euphorbia heterophylla* L.
 very common everywhere in openings and bare places.
- 3 75 *Enteromorpha*
 in bottom of intermittent ~~artificial~~ pool, varying with tides, but inland.
- 3 76 *Tineachne insularis* (L.) Nees
 very local, patches
- 3 77 *Cenchrus echinatus* L.
 common generally in open places.
- 3 78 *Fimbristylis cymosa* R.Br.
 common generally, especially in open places.
- 3 79 alga
 common on brackish pools in bomb craters.
- 4 80 *Cordia subcordata* Lam.
 subdominant ^{locally} in *Pisonia* forest

shrub 3 m. tall, spreading and tangled.

small tree, wood very soft; staminate flowers and fruits on same tree prostrate, extensive, forming mats; flowers purple.

dense tufts.

forming a dense floating mat.

tree 4 m. tall (others larger & trunks up to 6 dm or more thick). flowers bright orange.

- 34431 *Portulaca oleracea* L.
 2 common generally in
 open disturbed places
- 4 82 *Eleusine indica* (L.) Gaertn.
 common generally around
 buildings
- 3 83 *Lonchus deraeus* L.
 rare around dwellings.

varying between 1-3 m. tall,
 no large trunks seen.
 The *Pemphis* seems to grow
 on either fine silty sand
 or on broken coral rock.
 This belt varies in width
 up to 300 m. wide, but
 mostly much narrower,
 and is present pretty
 much around the lagoon
 shore of Wake Islet.
 On such short examination
 it is hard to evaluate the
 change in this area.

Over the ~~majority~~ ^{larger part} of
 the area of all three islets
 the vegetation was evidently
 almost completely destroyed.
 Ditches, tank traps, platforms,
 ruined buildings, gun
 emplacements, revetments,
 and underground instal-
 lations are almost every-
 where, with refuse of all

fleshy, spreading,
 purplish-green

small spreading leaf.

erect, flowers pale yellow.

sorts very abundant.
 According to Mursion, CAA
 manager, four years
 ago the vegetation, generally
 on the island, was about
 a foot high. Now the
 dominant growth is
 a stand of *Messerschmidia*
 10-15 or more feet high.
 Brown, engineer for CAA, says
 that in favorable sandy
 places *Messerschmidia*
 will grow 6 feet a year.
 Over large areas the present
 vegetation is a solid stand
 of *Messerschmidia*. On
 the south side, west of
 Peacock Pt., there is some
 admixture of *Scaevola*,
 not common elsewhere.
 A little *Pemphis* is scattered
 in here and there, generally.
 Near Peacock Point and
 elsewhere a scrub of

Cordia from 1-2 m. tall and very dense, is common locally. This is probably second-growth.

That the general *Messerschmidia* vegetation is second-growth is obvious from the lack of trunks more than 1 dm. thick and the fact that this scrub grows on embankments and piles of coral rubble that clearly date from the war.

In open places, generally, *Pluchea odorata* forms a scrub 1-2 m. tall. This is generally healthy, but in places is conspicuously chlorotic.

On denuded areas *Cenchrus echinatus* is common, and in many of them, *Euphorbia heterophylla* is abundant, at this season mostly completely dried up. Very chlorotic locally ^{where not dry}.

While Bryan noted only one patch of *Ipomoea pes-caprae* and one patch of *Gossypium*, both are locally common now on many parts of ~~the~~ Wake ~~Island~~ ^{and some other} ~~islands~~ and Peale Islet. ^{the latter esp. around the old airplane ramps.}

The Japanese must have planted the *Gossypium*, as it occurs consistently around their old installations.

In the intervals between the trees *Boerhavia* of two forms is extremely common, also an erect, green form of *Portulaca lutea*. *Portulaca oleracea* is commoner near installations.

Messerschmidia is said to be killed by salt water on its roots from very high tides - acc. Mr. Brown there is an area of 3 acres where it was killed and has not shown signs of coming back, as yet.

Some patches of *Cynodon dactylon* exist on the island. Much more was planted last fall, but few seeds sprouted, as there was little rain from December to April. *Messerschmidia* in many places looks dried up. *Pisonia* and *Cordia* are more or less defoliated, and the *Cordia* sometimes dead at the tips.

Here and there, in excavations and bomb craters, is standing ~~po~~ brackish water, one of the ponds being an acre or more in extent and locally called the Lake. These ponds and pools have a mat of ~~or~~ locally *Vaucheria*-like algae ^{locally} top, and flourishing populations of mosquito fish. There are also dragon-fly larvae. The water tastes perceptibly brackish. What so many fish live in, since the mosquito population has been reduced is not clear. The fish were planted about 8 months ago by Mr. Schultz, from a stock introduced from Hawaii about 4 years ago into a concrete reservoir on Peale I. by Suitono, a Samoan now with Pearl Harbor Police Dept. The fish are still numerous in this reservoir. There is one plant of *Eichhornia crassipes* left in this reservoir, ~~at~~ out of a formerly larger stand. They have been taken by people to plant in aquaria, etc.

A limnological study would be of interest.

Until a year ago there were said to be untold numbers of rats, flies, and mosquitoes (one *Culex*, two *Aedes* sp.) on the island.

Mr. Frederick Schultz.

3919 Waialae Ave. Honolulu 55, T.H.

(present % EAA, Wake I.)

in Feb. 1951 undertook pest control operations.

Mosquito breeding places were eliminated & stocked with mosquito fish. Garbage and other fly breeding material was eliminated, garbage dumps regularly burned. Rotted rats poisoned with Wolfen (1) was used in a systematic campaign against the rats. The success of this may be measured by the fact that in 2 1/2 days covering the island with Mr. Schultz one rat was seen, no mosquitoes, and only a very few flies. In his weekly examination of poisoned baits, in covered traps (like miniature quonset huts) he finds less than 5% touched by rats.

Feeding tests by Schultz showed that the rats commonly eat the young fruits and fleshy

fruiting bracts of *Ipomoea tuba*. He finds this quite consistent with the fact that rats were usually very numerous in tangles, and thickets, and mats of this plant. It would also supply water of which, there was none available naturally except after rains before the war.

The system of baiting was to establish parallel lines of baits across the island, ~~a~~ series of them, examining and re-baiting every day, till they were not disturbed much any more, then to move the series farther apart, doing the same thing over repeatedly, till the ends of the island were reached, then maintaining a few lines of baits as checks, examining these every week. As long as the % of baits visited is below 10% the rats are considered controlled.

Birds seen on Wake -

A few common noddies, flying, one nest but not sure if occupied.

A large rookery of sooty terns (? back and wings very black, crown black, forehead with a white ∇ , bill & feet black, under-parts white. - young sooty-brown, speckled with white on back, sooty gray beneath, when fully feathered. In pin feather stage light gray downy beneath, sooty brown on back and sides) on lagoon side of north hook of ~~Wake~~ Wake Islet. Young in various stages, no eggs seen.

Many frigate birds seen, principally roosting on remains of an old pier on Peale Islet in lagoon.

Occasional fairy terns flying.

Some boobies, nesting on Peale Islet, in trees. Sp. not det.

One tattler on lagoon beach.

Two plovers seen, but Mr. Schultze says there were many only a day earlier.

Dogs & cats both seen, domestic.

Gardens seem mostly unsuccessful when just planted in the ground. (see p. 130). Around an old Japanese garden site *Amaranthus viridis*, *A. tricolor* (?), *Ipomoea batatas* and *Nicotiana glauca* persist.

The method of gardening used most successfully is to frame up top-soil about 6" deep, surrounded by 1x6 boards. Then water this well, shade it with leafy branches of *Messerschmidia*. In this sort of arrangement, at Pranslean compound, tomatoes, lettuce, Broccoli, parsley, dill, onions, beans, ^{radishes}, were growing successfully.

Around this compound were small plants of *Ficus carica*, *Thespesia populnea*, *Ficus* sp., *Casuarina equisetifolia*, *Sorghum vulgare* var. *technicum* (broomcorn), *Catharanthus roseus*, *Terminalia catappa*, *Euphorbia pulcherrima*, *Cucurbita maxima* (flat white summer squash).

In pots, were *Codiaeum variegatum*, *Chrysophyllum cainito*?, *Bauhinia* sp., *Mangifera indica*, *Eucalyptus citriodora*, ~~*Lamanea*~~ *Guaiacum*?, *Capricorn putrescens*, *Zea mays* (reached maturity, but very spindly, with abortive ears, grown in a drum of soil. Cucumbers at least flowered. *Carica papaya* gets several feet tall, then dies.

Around houses in housing area - one coconut with trunk 2 m. tall; one *Pandanus tectorius* 2 m. tall.

In pots at least two species of *Kalanchoe*, *Bryophyllum pinnatum*, *Lansevieria zeylanicum* (?), *Echeveria* sp. etc.

The soils are generally coral rubble, with here and there a slight ~~area~~ coloring by humus, under *Pisonia* grows a black friable soil on the coral rubble. In one spot, site of old Japanese garden a small area of

dark brownish gray
fine soil. Part of this
is a cemented platform
of brown rock - possibly
phosphate. Sample #129.

The north side of
Wake Islet is a boulder
beach, with continuous
beach rock below high
tide level, overlain landward
by beach-conglomerate. This
is topped by the boulder beach
and boulder ridge.

This is the locality ~~so~~
described by Bryan 1927 as
having an impenetrable
barrier of vegetation.

Now it is completely
cleared, having only
scattered brush and
herbs. There were
large gun emplacements,
etc. here.

May 18 - Munson Hill
 s.e. of Falls Church
 old field being thickly
 invaded by mixed
 stand of young trees,
 some older trees.

34484 *Maclura pomifera*
 6 rare in

6 85 *Morus alba*
 occasional in.

5 86 *Acer saccharum* L.
 rare in grove on top of hill.

~~10 87 *Hedysotis purpurea* var. *purpurea*~~

May 25 Sleepy Hollow,
 1 1/2 mi. s. of Falls Church
 near Tripps Run

open slopes in pasture
 with scattered brush
 and springy ground

10 87 *Hedysotis purpurea* var. *purpurea*
 common on slope

5 88 ~~*Callitriche*~~ *Hydrocotyle ranunculoides* L.f.
 common in spring-fed pool.

5 89 *Hydrocotyle ranunculoides* L.f.
 common in spring-fed pool

small rounded tree
 5 m. tall, flowers ♂,
 mainly on older wood
 of long wand-like
 twigs.
 - shrub 2.5 m. tall,
 - fruits immature.
 - old tree 10 m. tall, sterile.

floating leaves expanded.

158

1952

Maryland

June 1 - Head of Broad Creek

shore of fine sand,
slightly brackish water

34490

Peltandra virginica

common

Anne Arundel Co.

159

clumps, leaf-blade
more or less erect,
quite variable from
clump to clump.

34490. (3)

34490-(3)

158

1952

Maryland

June 1 - Head of Broad Creek

shore of fine sand,
slightly brackish water34490
3*Peltandra virginica*
common

Anne Arundel Co.

159

clumps, leaf-blades
more or less erect,
quite variable from
clump to clump.

June 6 - Holmes Run,
just above Lake Barcroft
area cleared for sewer
10 years ago, with thickets
along creek banks, wooded

34491 *Fraxinus pennsylvanica* MARSH.

in bottom-land flooded
by low dam in creek

4 92 *Fraxinus americana* L.

same

4 93 *Betula nigra*

thickets along creek bank

6 94 *Physocarpus opulifolius*
thickets along creek bank

2 95 *Carex*

open grassy area

6 96 *Penstemon*

common in open area

bottom-land.

Tree 10 m. tall, partially
dead from flooding.

same

tree 10 m. tall, 2.5 dm.

thick, bark pinkish,

~~shaggy~~; ~~sterile~~

leaves glaucous

beneath; sterile.

shrub 2 m. tall, branches

arching; fruits dull

red.

culms weakly

arching.

erect; flowers

dull purplish white.

162

163

Dec. 16/52

Ngariungs Islet.

- ✓ 1 *Pisonia grandis* R. Br.
- ✓ 2 *Scaevola frutescens* (Mill.) Krause
- ✓ 3 *Messerschmidia argentea* (L.f.) Johnston.
- ✓ 4 *Barringtonia asiatica* (L.) Kurz
- ✓ 5 *Pisonia grandis* R. Br.
- ✓ 6 *Terminalia samoensis* Rech.
- ✓ 7 *Cassytha filiformis* L. (Kt. & Ghe.)
- ✓ 8 *Euphorbia chammissonis* Boiss.
- ✓ 9

tallest tree

tallest tree

parasitic on no 8

Dec. 16.

Ngajangel Islet (main islet)

- ✓ (2) 9 *Crataeva speciosa* Volk.
- ✓ 10 *Epipremnum pinnatum* (L.) Engl. &
- ✓ 11 *Vigna marina* (Burm.) Merr.
- ✓ 12 *Asplenium nidus* L.
- ✓ 13 *Nephrolepis hirsutula* (Forst.) Presl ?
- ✓ 14 *Ficus sp. rugifolia* Warb.
- ✓ 15 *Pipturus argenteus* (Forst.) Wedd.
- ✓ 16 *Epipremnum pinnatum* (L.) Engl. ?
- ✓ 17 *Ochromia oppositifolia* (Lam.) K. Schum.
- ✓ 18 *Pandanus tectorius* Pers.
- ✓ 19 *Acalypha amentacea* var. *carolinensis*
- ✓ 20 *Clerodendrum speciosissimum* Van Geert Volk.
- ✓ 21 *Cycas circinalis* (L.) Presl.
- ✓ 22 *Flemingia nuda* (L.) Gaertn. ex Wedd.
- ✓ 23 *Portulaca samoensis* v. Poelln.
- ✓ 24 ~~W. Arbygranthospora~~ *Canavalia microcarpa* (L.) Piper
- ✓ 25 *Elatosterna calcareum* Merr. ?
- ✓ 26 *Asystasia blumei* Nees
- ✓ 27 *Phyllanthus niruri* L.

young growth?

Tree 10 m. high

young growth?

Dec. 16

Ngajangel Islet.

- 28 Polyscias fruticosa (L.) Forst. Harms
 (L) 29 Eugenia javanica Lam.
 30 ~~30~~ Wedelia biflora (L.) DC.
 31 ~~Ficus~~ ~~sp~~ unrecognized seedling
 32 (L) ~~32~~ ~~Triumfetta~~ Triumfetta procumbens Forst.
 (L) 33 Achyranthes aspera L.
 34 (L) ~~34~~ ~~Cratogeomys~~ ~~repens~~ Holk. lepturus repens (Forst.) B. & H.
 35 Centella asiatica (L.) Urb.
 36 Polypodium scolopendria Burm. f.
 37 Nephrolepis ~~sp~~ biserrata (Sw.) Schott
 38 ~~38~~ ~~Adiantum~~ ~~oppositifolium~~ Holk. ~~Adiantum~~ ~~oppositifolium~~ + Ochonia oppositifolia (Lam.) K. Schum. ?
 39 Acalypha amataea var. carolinensis
 40 ~~40~~ ~~Digitaria~~ Digitaria longissima Mey
 41 Ischaemum muticum L.
 42 Eragrostis amabilis (L.) W. & A.
 43a Digitaria pruriens var
 44a ~~Lepturus~~ ~~lepturus~~ L.
 45a ~~lepturus~~
 46a ~~lepturus~~ ~~lepturus~~
 47a ~~Triumfetta~~ ~~procumbens~~
 48a ~~lepturus~~ ~~repens~~
 49a
 50a
 51a
 52a ~~Riptocarpus~~ ~~argenteus~~
 53a
 54a
 55a
 56a
 43b Digitaria timorensis (Kunth) Balansa

opposite
seedling

Dec. 1952

Auluptagel Island near Koror.

44 ~~524~~ ⁽²⁾ *Ponapea palauensis* Kaneh.5045 ~~524~~ ⁽²⁾ *Gulubiospis palauensis* Kaneh.46 ~~524~~ *Pandanus aimirihensis* Mart.

G. C. Gessitt coll.

also common in hill jungle on Babelthwap

172

173

# 112	-	p. 24	111 - 12
113	-	p. 24	
114	-	p. 56	
115	-	p. 58	
116	-	p. 60	
117	-	p. 60	
118	-	p. 61	
119	-	p. 61	
120	-	p. 59	
121	-	p. 66	
122	-	p. 66	
123	-	p. 66	
124	-	p. 66	
125	-	p. 66	
126	-	p. 70	
127	-	p. 74	
128	-	p. 73	
129	-	p. 152	

Index to Vials

- # 133 - p. 24
- 134 - p. 24
- 135 - p. 56
- 136 - p. 58
- 137 - p. 70
- 138 - p. 71
- 139 - p. 65
- 140 - p. 116
- 141 - p. 132
- 142 - p. 132
- 143 - p. 132
- 144 - p. 132
- 145 - p. 132

19 200



